arXiv:1406.3942v1 [math.GN] 16 Jun 2014

Towards a Descriptive Theory of cb_0-Spaces

Victor Selivanov
A.P. Ershov Institute of Informatics Systems SB RAS
Novosibirsk, Russia

June 17, 2014

Abstract

The paper tries to extend results of the classical Descriptive Set Theory to as many countably based T_0-spaces (cb_0-spaces) as possible. Along with extending some central facts about Borel, Luzin and Hausdorff hierarchies of sets we consider also the more general case of k-partitions. In particular, we investigate the difference hierarchy of k-partitions and the fine hierarchy closely related to the Wadge hierarchy.

Key words. Borel hierarchy, Luzin hierarchy, Hausdorff hierarchy, Wadge reducibility, cb_0-space, k-partition, h-preorder, well preorder.

1 Introduction

Classical Descriptive Set Theory [Ke95] is an important field of mathematics with numerous applications. It investigates descriptive complexity of sets, functions and equivalence relations in Polish (i.e., separable complete metrizable) spaces.

Although Polish spaces are sufficient for many fields of classical mathematics, they are certainly not sufficient for many fields of Theoretical Computer Science where non-Hausdorff spaces (in particular, ω-continuous domains) and non-countably-based spaces (in particular, Kleene-Kreisel continuous functionals) are of central importance. For this reason, the task of extending classical Descriptive Set Theory (DST) to as many non-Polish spaces as possible attracted attention of several researchers.

Some parts of DST for ω-continuous domains (which are typically non-Polish) were developed in [Se04, Se05a, Se06, Se08]. In [Br13] a good deal of DST was developed for the so called quasi-Polish spaces (see the next section for a definition of this class of cb_0-spaces) which include both the Polish spaces and the ω-continuous domains. For some attempts to develop DST for non-countably based spaces see e.g. [JR82, MS10, FHK11, P12, PdB13].

In this paper, we try to develop DST for some classes of cb_0-spaces beyond the class of quasi-Polish spaces. As is usual in classical DST, we put emphasis on the "ininitary version" of hierarchy theory where people are concerned with transfinite (along with finite) levels of hierarchies. The "finitary" version where people concentrate on the finite levels of hierarchies has a special flavor and is relevant to several fields of Logic and Computation Theory; it was systematized in [Se06, Se08a, Se12].

We extend some well known facts about classical hierarchies in Polish spaces to natural classes of cb_0-spaces. Namely, we show that some levels of hierarchies of cb_0-spaces introduced in [ScSe13, ScSe14] provide natural examples of classes of cb_0-spaces with reasonable DST (in particular, the classical Suslin, Hausdorff-Kuratowski and non-collapse theorems for the Borel, Luzin and Hausdorff hierarchies are true for such spaces). This portion of our results are technically easy and follow rather straightforwardly from the classical DST and some notions and results in [MSS12, ScSe13].

Along with the classical hierarchies of sets we are interested also in the difference and fine hierarchies of k-partitions [Se06, Se07, Se07a, Se08, Se08a, Se11] which seem to be natural, non-trivial and useful generalization of the corresponding hierarchies of sets. Also, along with the classical Wadge reducibility [Wad72, Wad84, VW76] we discuss its extension to k-partitions [Her93, Her96, Se07a], and some their weaker versions introduced and studied in [AM03, And06, MR09, MSS12].

Already the extension of the Hausdorff difference hierarchy to k-partitions is a non-trivial task. The general "right" finitary version of this hierarchy was found only recently in [Se12], although for some

particular cases it was already in our previous publications. The general "right" infinitary version of this hierarchy is new here, although the definition adequate for bases with the ω-reduction property was also found earlier [Se07, Se07a, Se08]. That the definition in this paper is right follows from the nice properties of the difference hierarchy of k-partition (in particular, the natural version of the Hausdorff-Kuratowski theorem).
The situation with the fine hierarchy (which aims to extend the Wadge hierarchy to the case of sets and k-partitions in arbitrary spaces) is even more complicated. This task is not obvious even for the case of sets because the Wadge hierarchy is developed so far only for the Baire space (and some of its close relatives) in terms of m-reducibility by continuous functions and with a heavy use of Martin determinacy theorem [Wad72, Wad84, VW76]. As a result, there is no clear explicit description of levels of the hierarchy in terms of set-theoretic operations which one could try to extend to other spaces (more precisely, some rather indirect descriptions presented in [Wad84, Lo83] strongly depend on the ω-reduction property of the open sets which usually fails in non-zero-dimensional spaces). Probably, that was the reason why some authors tried to obtain alternative characterizations of levels of the Wadge hierarchy [Lo83, Du01]. In a series of our papers (see e.g. [Se06, Se08a]) a characterization of levels of an abstract version of Wadge hierarchy in the finitary case was achieved that was extended in [Se12] to the case of k-partitions. Here we develop an infinitary version of this approach and try to explain why the corresponding hierarchy is the "right" extension of the Wadge hierarchy to arbitrary spaces and to the k-partitions. Since the notation and full proofs in this context are extremely involved, we concentrate here only on the partitions of finite Borel rank and avoid the complete proofs of some complicated results, giving only precise formulations and hints of proofs with references to closely related earlier proofs in the finitary context.
After recalling some notions and known facts in the next section, we discuss some basic properties of Borel and Luzin hierarchies in cb_0-spaces in Section 3. In Section 4 we establish some basic facts on the difference hierarchies of k-partitions in cb_0-spaces. The main result here is the Hausdorff-Kuratowski theorem for k-partitions in quasi-Polish spaces. In Section 5 we extend the difference hierarchies of k-partitions to the the fine hierarchies of k-partitions. In particular, we extend the Hausdorff-Kuratowski theorem to the fine hierarchy. We conclude in Section 6 with sketching a possible further research on extending the classical Descriptive Set Theory.

2 Notation and preliminaries

In this section we recall some notation, notions and results used in the subsequent sections.

2.1 cb_0-Spaces and qcb_0-spaces

Here we recall some topological notions and facts relevant to this paper.
We freely use the standard set-theoretic notation like $dom(f)$, $rng(f)$ and $graph(f)$ for the domain, range and graph of a function f, respectively, $X \times Y$ for the Cartesian product, and $P(X)$ for the set of all subsets of X. For $A \subseteq X$, $\overline{A}$ denotes the complement $X \setminus A$ of A in X. We identify the set of natural numbers with the first infinite ordinal ω. The first uncountable ordinal is denoted by ω_1. The notation $f : X \to Y$ means that f is a total function from a set X to a set Y.
We assume the reader to be familiar with the basic notions of topology [En89]. The collection of all open subsets of a topological space X (i.e. the topology of X) is denoted by $\mathcal{O}(X)$; for the underlying set of X we will write X in abuse of notation. We will often abbreviate "topological space" to "space". A space is *zero-dimensional* if it has a basis of clopen sets. Recall that a *basis* for the topology on X is a set $\mathcal{B}$ of open subsets of X such that for every $x \in X$ and open U containing x there is $B \in \mathcal{B}$ satisfying $x \in B \subseteq U$.
Let ω be the space of non-negative integers with the discrete topology. Of course, the spaces $\omega \times \omega = \omega^2$, and $\omega \sqcup \omega$ are homeomorphic to ω, the first homeomorphism is realized by the Cantor pairing function $\langle \cdot, \cdot \rangle$.
Let $\mathcal{N} = \omega^\omega$ be the set of all infinite sequences of natural numbers (i.e., of all functions $\xi \colon \omega \to \omega$). Let ω^* be the set of finite sequences of elements of ω, including the empty sequence. For $\sigma \in \omega^*$ and $\xi \in \mathcal{N}$, we write $\sigma \sqsubseteq \xi$ to denote that σ is an initial segment of the sequence ξ. By $\sigma\xi = \sigma \cdot \xi$ we denote the concatenation of σ and ξ, and by $\sigma \cdot \mathcal{N}$ the set of all extensions of σ in $\mathcal{N}$. For $x \in \mathcal{N}$, we can write

$x = x(0)x(1)\dots$ where $x(i) \in \omega$ for each $i < \omega$. For $x \in \mathcal{N}$ and $n < \omega$, let $x^{<n} = x(0)\dots x(n-1)$ denote the initial segment of x of length n. Notations in the style of regular expressions like 0^ω, 0^*1 or 0^m1^n have the obvious standard meaning.
By endowing $\mathcal{N}$ with the product of the discrete topologies on ω, we obtain the so-called *Baire space*. The product topology coincides with the topology generated by the collection of sets of the form $\sigma \cdot \mathcal{N}$ for $\sigma \in \omega^*$. The Baire space is of primary importance for Descriptive Set Theory and Computable Analysis. The importance stems from the fact that many countable objects are coded straightforwardly by elements of $\mathcal{N}$, and it has very specific topological properties. In particular, it is a perfect zero-dimensional space and the spaces $\mathcal{N}^2, \mathcal{N}^\omega, \omega \times \mathcal{N} = \mathcal{N} \sqcup \mathcal{N} \sqcup \cdots$ (endowed with the product topology) are all homeomorphic to $\mathcal{N}$. Let $(x, y) \mapsto \langle x, y\rangle$ be a homeomorphism between $\mathcal{N}^2$ and $\mathcal{N}$. The subspace $\mathcal{C} := 2^\omega$ of $\mathcal{N}$ formed by the infinite binary strings (endowed with the relative topology inherited from $\mathcal{N}$) is known as the *Cantor space*.
The *Sierpinski space* $\mathbb{S}$ is the two-point set $\{\bot, \top\}$ where the set $\{\top\}$ is open but not closed. The space $P\omega$ is formed by the set of subsets of ω equipped with the Scott topology. A countable base of the Scott topology is formed by the sets $\{A \subseteq \omega \mid F \subseteq A\}$, where F ranges over the finite subsets of ω. Note that $P\omega = \mathcal{O}(\omega)$. As is well-known [GH80], $P\omega$ is universal for cb_0-spaces:

Proposition 2.1 *A topological space X embeds into $P\omega$ iff X is a cb_0-space.*

Remember that a space X is *Polish* if it is countably based and metrizable with a metric d such that (X, d) is a complete metric space. Important examples of Polish spaces are ω, $\mathcal{N}$, $\mathcal{C}$, the space of reals $\mathbb{R}$ and its Cartesian powers $\mathbb{R}^n$ $(n < \omega)$, the closed unit interval $[0, 1]$, the Hilbert cube $[0, 1]^\omega$ and the Hilbert space $\mathbb{R}^\omega$. Simple examples of non-Polish spaces are $\mathbb{S}$, $P\omega$ and the space $\mathbb{Q}$ of rationals.
A space X is *quasi-Polish* [Br13] if it is countably based and quasi-metrizable with a quasi-metric d such that (X, d) is a complete quasi-metric space. A *quasi-metric* on X is a function from $X \times X$ to the nonnegative reals such that $d(x, y) = d(y, x) = 0$ iff $x = y$, and $d(x, y) \le d(x, z) + d(z, y)$. Since for the quasi-metric spaces different notions of completeness and of a Cauchy sequence are considered, the definition of quasi-Polish spaces should be made more precise (see [Br13] for additional details). We skip these details because we will in fact use another characterization of these spaces given below. Note that the spaces $\mathbb{S}$, $P\omega$ are quasi-Polish while the space $\mathbb{Q}$ is not.
A *representation* of a space X is a surjection of a subspace of the Baire space $\mathcal{N}$ onto X. A basic notion of Computable Analysis is the notion of admissible representation. A representation δ of X is *admissible*, if it is continuous and any continuous function $\nu : Z \to X$ from a subset $Z \subseteq \mathcal{N}$ to X is continuously reducible to δ, i.e. $\nu = \delta \circ g$ for some continuous function $g : Z \to \mathcal{N}$. A topological space is *admissibly representable*, if it has an admissible representation.
The notion of admissibility was introduced in [KW85] for representations of cb_0-spaces (in a different but equivalent formulation) and was extensively studied by many authors. In [Sch02, Sch03] the notion was extended to non-countably based spaces and a nice characterization of the admissibly represented spaces was achieved. Namely, the admissibly represented sequential topological spaces coincide with the qcb_0-spaces, i.e., T_0-spaces which are topological quotients of countably based spaces.
In [Br13] the following important characterization of quasi-Polish spaces in terms of Borel hierarchy was obtained.

Proposition 2.2 *A space is quasi-Polish iff it is homeomorphic to a $\mathbf{\Pi}^0_2$-subset of $P\omega$ with the induced topology.*

2.2 Hierarchies of sets

Here we briefly recall definitions and some properties of Borel and Luzin hierarchies in arbitrary topological spaces.
A *pointclass* in a space X is simply a collection $\mathbf{\Gamma}(X)$ of subsets of X. A *family of pointclasses* [Se13] is a family $\mathbf{\Gamma} = \{\mathbf{\Gamma}(X)\}$ indexed by arbitrary topological spaces X such that each $\mathbf{\Gamma}(X)$ is a pointclass on X and $\mathbf{\Gamma}$ is closed under continuous preimages, i.e. $f^{-1}(A) \in \mathbf{\Gamma}(X)$ for every $A \in \mathbf{\Gamma}(Y)$ and every continuous function $f\colon X \to Y$. A basic example of a family of pointclasses is given by the family $\mathcal{O} = \{\mathcal{O}(X)\}$ of the topologies of all the spaces X.

We will use some operations on families of pointclasses. First, the usual set-theoretic operations will be applied to the families of pointclasses pointwise: for example, the union $\bigcup_i \mathbf{\Gamma}_i$ of the families of pointclasses $\mathbf{\Gamma}_0, \mathbf{\Gamma}_1, \dots$ is defined by $(\bigcup_i \mathbf{\Gamma}_i)(X) = \bigcup_i \mathbf{\Gamma}_i(X)$.
Second, a large class of such operations is induced by the set-theoretic operations of L.V. Kantorovich and E.M. Livenson (see e.g. [Se13] for the general definition). Among them are the operation $\mathbf{\Gamma} \mapsto \mathbf{\Gamma}_\sigma$, where $\mathbf{\Gamma}(X)_\sigma$ is the set of all countable unions of sets in $\mathbf{\Gamma}(X)$, the operation $\mathbf{\Gamma} \mapsto \mathbf{\Gamma}_\delta$, where $\mathbf{\Gamma}(X)_\delta$ is the set of all countable intersections of sets in $\mathbf{\Gamma}(X)$, the operation $\mathbf{\Gamma} \mapsto \mathbf{\Gamma}_c = \check{\mathbf{\Gamma}}$, where $\mathbf{\Gamma}(X)_c$ is the set of all complements of sets in $\mathbf{\Gamma}(X)$, the operation $\mathbf{\Gamma} \mapsto \mathbf{\Gamma}_d$, where $\mathbf{\Gamma}(X)_d$ is the set of all differences of sets in $\mathbf{\Gamma}(X)$, the operation $\mathbf{\Gamma} \mapsto \mathbf{\Gamma}_\exists$ defined by $\mathbf{\Gamma}_\exists(X) := \{\exists^{\mathcal{N}}(A) \mid A \in \mathbf{\Gamma}(\mathcal{N} \times X)\}$, where $\exists^{\mathcal{N}}(A) := \{x \in X \mid \exists p \in \mathcal{N}.(p,x) \in A\}$ is the projection of $A \subseteq \mathcal{N} \times X$ along the axis $\mathcal{N}$, and finally the operation $\mathbf{\Gamma} \mapsto \mathbf{\Gamma}_\forall$ defined by $\mathbf{\Gamma}_\forall(X) := \{\forall^{\mathcal{N}}(A) \mid A \in \mathbf{\Gamma}(\mathcal{N} \times X)\}$, where $\forall^{\mathcal{N}}(A) := \{x \in X \mid \forall p \in \mathcal{N}.(p,x) \in A\}$.
The operations on families of pointclasses enable to provide short uniform descriptions of the classical hierarchies in arbitrary spaces. E.g., the Borel hierarchy is the family of pointclasses $\{\mathbf{\Sigma}^0_\alpha\}_{\alpha<\omega_1}$ defined by induction on α as follows [Se06, Br13]: $\mathbf{\Sigma}^0_0(X) := \{\emptyset\}$, $\mathbf{\Sigma}^0_1 := \mathcal{O}$, $\mathbf{\Sigma}^0_2 := (\mathbf{\Sigma}^0_1)_{d\sigma}$, and $\mathbf{\Sigma}^0_\alpha(X) := (\bigcup_{\beta<\alpha} \mathbf{\Sigma}^0_\beta(X))_{c\sigma}$ for $\alpha > 2$. The sequence $\{\mathbf{\Sigma}^0_\alpha(X)\}_{\alpha<\omega_1}$ is called *the Borel hierarchy* in X. We also let $\mathbf{\Pi}^0_\beta(X) := (\mathbf{\Sigma}^0_\beta(X))_c$ and $\mathbf{\Delta}^0_\alpha(X) := \mathbf{\Sigma}^0_\alpha(X) \cap \mathbf{\Pi}^0_\alpha(X)$. The classes $\mathbf{\Sigma}^0_\alpha(X), \mathbf{\Pi}^0_\alpha(X), \mathbf{\Delta}^0_\alpha(X)$ are called the *levels* of the Borel hierarchy in X.
We recall an important structural property of $\mathbf{\Sigma}$-levels of the Borel hierarchy. Let Γ be a family of pointclasses. A pointclass $\Gamma(X)$ has the *ω-reduction property* if for each countable sequence $A_0, A_1, \dots$ in $\Gamma(X)$ there is a countable sequence $D_0, D_1, \dots$ in $\Gamma(X)$ such that $D_i \subseteq A_i$, $D_i \cap D_j = \emptyset$ for all $i \neq j$ and $\bigcup_{i<\omega} D_i = \bigcup_{i<\omega} A_i$.

Proposition 2.3 *For any space X and any $2 \leq \alpha < \omega_1$, $\mathbf{\Sigma}^0_\alpha(X)$ has the ω-reduction properties. If X is zero-dimensional, the same holds for the class $\mathbf{\Sigma}^0_1(X)$ of open sets.*

The *hyperprojective hierarchy* is the family of pointclasses $\{\mathbf{\Sigma}^1_\alpha\}_{\alpha<\omega_1}$ defined by induction on α as follows: $\mathbf{\Sigma}^1_0 = \mathbf{\Sigma}^0_2$, $\mathbf{\Sigma}^1_{\alpha+1} = (\mathbf{\Sigma}^1_\alpha)_{c\exists}$, $\mathbf{\Sigma}^1_\lambda = (\mathbf{\Sigma}^1_{<\lambda})_{\delta\exists}$, where $\alpha, \lambda < \omega_1$, λ is a limit ordinal, and $\mathbf{\Sigma}^1_{<\lambda}(X) := \bigcup_{\alpha<\lambda} \mathbf{\Sigma}^1_\alpha(X)$.
In this way, we obtain for any topological space X the sequence $\{\mathbf{\Sigma}^1_\alpha(X)\}_{\alpha<\omega_1}$, which we call here *the hyperprojective hierarchy in X*. The pointclasses $\mathbf{\Sigma}^1_\alpha(X)$, $\mathbf{\Pi}^1_\alpha(X) := (\mathbf{\Sigma}^1_\alpha(X))_c$ and $\mathbf{\Delta}^1_\alpha(X) := \mathbf{\Sigma}^1_\alpha(X) \cap \mathbf{\Pi}^1_\alpha(X)$ are called *levels of the hyperprojective hierarchy in X*. The finite non-zero levels of the hyperprojective hierarchy coincide with the corresponding levels of the Luzin's projective hierarchy [Br13, ScSe13]. The class of *hyperprojective sets* in X is defined as the union of all levels of the hyperprojective hierarchy in X. For more information on the hyperprojective hierarchy see [Ke83, Ke95, ScSe14]. Below we will also consider some other hierarchies, in particular the Hausdorff difference hierarchy.

2.3 k-Partitions and hierarchies over well posets

Here we discuss a more general notion of a hierarchy (compared with the notion of hierarchy of sets [Se08a, Se12]) which applies, in particular, to the hierarchies of k-partitions.
Let $2 \leq k < \omega$. By a *k-partition of a space X* we mean a function $A : X \to k = \{0, \dots, k-1\}$ often identified with the sequence $(A_0, \dots, A_{k-1})$ where $A_i = A^{-1}(i)$. Obviously, 2-partitions of X are identified with the subsets of X using the characteristic functions. The set of all k-partitions of X is denoted k^X. For $\mathbf{\Gamma} \subseteq P(X)$, let $(\mathbf{\Gamma})_k$ denote the set of k-partitions $A \in k^X$ such that $A_0, \dots, A_{k-1} \in \mathbf{\Gamma}(X)$. In particular, $(\mathbf{\Sigma}_{<\omega}(X))_k$ is the set of k-partitions of finite Borel rank which will be considered in Section 5.2.
The Wadge reducibility on subsets of X is naturally extended on k-partitions: for $A, B \in k^X$, $A \leq_W B$ means that $A = B \circ f$ for some continuous function f on X, In this way, we obtain the preorder $(k^X; \leq_W)$ which for $k \geq 3$ turns out much more complicated than the structure of Wadge degrees, even for simple case $X = \mathcal{N}$ [Her93, Her96, Se06, Se11].
To find the "right" extensions of the classical difference and Wadge hierarchies from the case of sets to the case of k-partitions is a quite non-trivial task. A reason is that levels of hierarchies of sets are always semi-well-ordered by inclusion (in particular, there are no three levels which are pairwise incomparable by inclusion) while the structure of hierarchies of k-partitions for $k \geq 3$ is usually more complicated than the structure of the hierarchies of sets (in particular, for $k \geq 3$ the poset of levels of difference hierarchies

of k-partitions under inclusion usually has antichains with any finite number of elements).
Here we recall from [Se12] a very general notion of a hierarchy that covers all hierarchies we discuss in this paper.

Definition 2.4 (1) For any poset P and any set A, by a P-hierarchy in A we mean a family $\{H_p\}_{p\in P}$ of subsets of A such that $p \leq q$ implies $H_p \subseteq H_q$.
(2) Levels (resp. constituents) of a P-hierarchy $\{H_p\}$ are the sets $H_{p_0} \cap \dots \cap H_{p_n}$ (resp. the sets $C_{p_0,\dots,p_n} = (H_{p_0} \cap \dots \cap H_{p_n}) \setminus \bigcup\{H_q \mid q \in P \setminus \uparrow \{p_0,\dots,p_n\}\}$) where $n \geq 0$ and $\{p_0,\dots,p_n\}$ is an antichain in P.
(3) A P-hierarchy $\{H_p\}$ is precise if $p \leq q$ is equivalent to $H_p \subseteq H_q$.

Note that the classical hierarchies of sets are obtained from the above definition if $A = P(X)$ and $P = \bar{2}\cdot\eta$ is the poset obtained by replacing any element of the ordinal η by an antichain with two elements, and that the notion of preciseness extends the non-collapse property of hierarchies. Note that levels of the classical hierarchies coincide with levels in the sense of the definition above. The constituents of say, Borel hierarchy, are $\mathbf{\Sigma}^0_\alpha \setminus \mathbf{\Pi}^0_\alpha$, $\mathbf{\Pi}^0_\alpha \setminus \mathbf{\Sigma}^0_\alpha$, $\mathbf{\Delta}^0_{\alpha+1} \setminus (\mathbf{\Sigma}^0_\alpha \cup \mathbf{\Pi}^0_\alpha)$, and $\mathbf{\Delta}^0_\lambda \setminus \bigcup_{\alpha<\lambda} \mathbf{\Sigma}^0_\alpha$, where λ is a limit countable ordinal.
As it was already mentioned, for hierarchies of k-partitions (obtained when $A = k^X$) we cannot hope to deal only with semi-well-ordered posets $P = \bar{2}\cdot\eta$ in the definition above. Fortunately, a slight weakening of this property is sufficient for our purposes: we can confine ourselves with the so called well posets (wpo) or, more generally well preorders (wqo). Recall that a wqo is a preorder P that has neither infinite descending chains nor infinite antichains. The theory of wqo (widely known as the wqo-theory) is a well developed field with several deep results and applications, see e.g. [Kru72]. It is also of great interest to hierarchy theory. An important role in wqo-theory belongs to the rather technical notion of a better preorder (bqo). Bqo's form a subclass of wqo's with good closure properties.
Note that if P is a wpo then the structure $(\{H_p \mid p \in P\}; \subseteq)$ of levels of a P-hierarchy under inclusion is also a wpo, hence some important features of the hierarchies of sets hold also for the hierarchies of partitions. Moreover, for such hierarchies we have some important properties of constituents, in particular the constituents form a partition of the set $\bigcup\{H_p \mid p \in P\}$ (see also Section 7 of [Se12] for additional details).
Although well posets are very simple compared with arbitrary posets, they are much more complicated than the semi-well-orders which essentially reduce to the ordinals. Obviously, there are a lot isomorphism types of well posets of a fixed rank. Below we consider some examples of well posets suitable for naming the levels of the difference and fine hierarchies of k-partitions.

2.4 Hierarchies of cb_0-spaces and qcb_0-spaces

Here we recall some classifications of qcb_0-spaces induced by the classical hierarchies of sets.
For any representation δ of a space X, let $EQ(\delta) := \{\langle p,q\rangle \in \mathcal{N} \mid p,q \in dom(\delta) \wedge \delta(p) = \delta(q)\}$. Let $\mathbf{\Gamma}$ be a family of pointclasses. A qcb_0-space X is called $\mathbf{\Gamma}$-*representable*, if X has an admissible representation δ with $EQ(\delta) \in \mathbf{\Gamma}(\mathcal{N})$. The class of all $\mathbf{\Gamma}$-representable spaces is denoted $\mathsf{QCB_0}(\mathbf{\Gamma})$. A cb_0-space X is called a $\mathbf{\Gamma}$-*space*, if X is homeomorphic to a $\mathbf{\Gamma}$-subspace of $P\omega$. The class of all $\mathbf{\Gamma}$-spaces is denoted $\mathsf{CB_0}(\mathbf{\Gamma})$.
These notions from [ScSe13] enable to transfer hierarchies of sets to the corresponding hierarchies of qcb_0-spaces. In particular, we arrive at the following definition.

Definition 2.5 The sequence $\{\mathsf{CB_0}(\mathbf{\Sigma}^0_\alpha)\}_{\alpha<\omega_1}$ (resp. the sequence $\{\mathsf{QCB_0}(\mathbf{\Sigma}^0_\alpha)\}_{\alpha<\omega_1}$) is called the *Borel hierarchy* of cb_0-spaces (resp. of qcb_0-spaces). By *levels* of this hierarchy we mean the classes $\mathsf{CB_0}(\mathbf{\Sigma}^0_\alpha)$ as well as the classes $\mathsf{CB_0}(\mathbf{\Pi}^0_\alpha)$ and $\mathsf{CB_0}(\mathbf{\Delta}^0_\alpha)$. In a similar way one can define the hyperprojective hierarchies of cb_0- and of qcb_0-spaces.

The following fact from [ScSe13, ScSe14] shows that the introduced hierarchies agree on cb_0-spaces:

Proposition 2.6 *For any* $\mathbf{\Gamma} \in \{\mathbf{\Pi}^0_2, \mathbf{\Sigma}^0_\beta, \mathbf{\Pi}^0_\beta, \mathbf{\Sigma}^1_\alpha, \mathbf{\Pi}^1_\alpha \mid 1 \leq \alpha < \omega_1, 3 \leq \beta < \omega_1\}$, *we have* $\mathsf{QCB_0}(\mathbf{\Gamma}) \cap \mathsf{CB_0} = \mathsf{CB_0}(\mathbf{\Gamma})$, *where* $\mathsf{CB_0}$ *is the class of all* cb_0*-spaces.*

Note that, by Proposition 2.2, $\mathsf{CB_0}(\mathbf{\Pi}^0_2)$ coincides with the class of quasi-Polish spaces.

3 Borel and Luzin hierarchies

In this section we extend some classical facts on the Borel and Luzin hierarchies in Polish spaces on larger classes of cb_0-spaces.

3.1 Some reducibilities and isomorphisms

Here we provide some information on versions of the Wadge reducibility and of the notion of homeomorphisms relevant to this paper.
Let $\mathbf{\Gamma}$ be a family of pointclasses and X, Y be topological spaces. By $\mathbf{\Gamma}(X, Y)$ we denote the class of functions $f : X \to Y$ such that $f^{-1}(A) \in \mathbf{\Gamma}(X)$ whenever $A \in \mathbf{\Gamma}(Y)$. A set $A \subseteq X$ is $\mathbf{\Gamma}$-*reducible* to a set $B \subseteq X$ (in symbols, $A \leq_{\mathbf{\Gamma}} B$) if $A = f^{-1}(B)$ for some $f \in \mathbf{\Gamma}(X, X)$.
Note that the $\mathbf{\Sigma}^0_1$-functions coincide with the continuous functions and the $\mathbf{\Sigma}^0_1$-reducibility coincides with the classical Wadge reducibility. $\mathbf{\Sigma}^0_\alpha$-Functions and $\mathbf{\Sigma}^0_\alpha$-reducibilities were investigated in [AM03, And06, MR09].
We say that topological spaces X, Y are $\mathbf{\Gamma}$-*isomorphic* if there is a bijection f between X and Y such that $f \in \mathbf{\Gamma}(X, Y)$ and $f^{-1} \in \mathbf{\Gamma}(Y, X)$. It is a classical fact of Descriptive Set Theory that every two uncountable Polish spaces X, Y are $\mathbf{\Delta}^1_1$-isomorphic (see e.g. [Ke95, Theorem 15.6]). The next result from [MSS12] extends this fact to the context of uncountable quasi-Polish spaces and computes an upper bound for the complexity of the Borel-isomorphism.

Proposition 3.1 *Let X, Y be two uncountable quasi-Polish spaces. Then X and Y are $\mathbf{\Delta}^0_{<\omega}$-isomorphic. If the inductive dimensions $dim(X)$, $dim(Y)$ of X, Y are distinct from ∞ then X and Y are $\mathbf{\Delta}^0_3$-isomorphic.*

Let again $\mathbf{\Gamma}$ be a family of pointclasses. By a $\mathbf{\Gamma}$-*family of pointclasses* we mean a family $\{E(X)\}_X$ indexed by arbitrary spaces such that $E(X)$ is a pointset in X, and $f^{-1}(A) \in E(X)$ for all $A \in E(Y)$ and $f \in \mathbf{\Gamma}(X, Y)$. Obviously, the $\mathbf{\Sigma}^0_1$-families of pointclasses are precisely the "usual" families of pointclasses.

Lemma 3.2 *Let $\mathbf{\Gamma}$ be a family of pointclasses. Then $\mathbf{\Gamma}$ is a $\mathbf{\Gamma}$-family of pointclasses, any continuous function $f : X \to Y$ is in $\mathbf{\Gamma}(X, Y)$, and any $\mathbf{\Gamma}$-family of pointclasses is a family of pointclasses.*

Proof. The first assertion is obvious. For the second assertion, let $f : X \to Y$ be continuous and $A \in \mathbf{\Gamma}(Y)$. Since $\mathbf{\Gamma}$ is a family of pointclasses, $f^{-1}(A) \in \mathbf{\Gamma}(X)$. Since A was arbitrary, $f \in \mathbf{\Gamma}(X, Y)$. The third assertion follows from the second one. □

Lemma 3.3 *Let $\alpha < \beta < \omega_1$. Then any $\mathbf{\Sigma}^0_\alpha$-function is a $\mathbf{\Sigma}^0_\beta$-function, and any $\mathbf{\Sigma}^1_\alpha$-function is a $\mathbf{\Sigma}^1_\beta$-function.*

Proof Hint. Proof is straightforward by induction on β, so we consider only the first assertion for the case $\beta = \alpha + 1$, as an example. Let $A \in \mathbf{\Sigma}^0_\beta(Y)$ and $f \in \mathbf{\Sigma}^0_\alpha(X, Y)$. Then $A = \bigcup_n (Y \setminus A_n)$ for some $A_0, A_1, \ldots \in \mathbf{\Sigma}^0_\alpha(Y)$. Then $f^{-1}(A) = \bigcup_n (X \setminus f^{-1}(A_n)) \in \mathbf{\Sigma}^0_\beta(X)$, so f is a $\mathbf{\Sigma}^0_\beta$-function. □

Proposition 3.4 (1) *Let $\mathbf{\Gamma} \in \{\mathbf{\Sigma}^0_\alpha, \mathbf{\Pi}^0_\alpha, \mathbf{\Sigma}^1_\beta, \mathbf{\Pi}^1_\beta \mid \omega \leq \alpha < \omega_1, 1 \leq \beta < \omega_1\}$ and $X \in \mathsf{CB}_0(\mathbf{\Gamma})$. Then X is $\mathbf{\Gamma}$-isomorphic to a subspace S of $\mathcal{N}$ such that $S \in \mathbf{\Gamma}(\mathcal{N})$.*

(2) *Let $\mathbf{\Gamma} \in \{\mathbf{\Sigma}^0_\alpha, \mathbf{\Pi}^0_\alpha, \mathbf{\Sigma}^1_\beta, \mathbf{\Pi}^1_\beta \mid 3 \leq \alpha < \omega_1, 1 \leq \beta < \omega_1\}$ and $X \in \mathsf{CB}_0(\mathbf{\Gamma})$, $dim(X) \neq \infty$. Then X is $\mathbf{\Gamma}$-isomorphic to a subspace S of $\mathcal{N}$ such that $S \in \mathbf{\Gamma}(\mathcal{N})$.*

Proof. Both items are checked in the same way, so consider only the first one. Assume without loss of generality that $X \in \mathbf{\Gamma}(P\omega)$. By Proposition 3.1, the spaces $P\omega$ and $\mathcal{N}$ are $\mathbf{\Delta}^0_{<\omega}$-isomorphic, hence also $\mathbf{\Sigma}^0_\omega$-isomorphic. By Lemma 3.3 $P\omega$ and $\mathcal{N}$ are $\mathbf{\Gamma}$-isomorphic, let $f : P\omega \to \mathcal{N}$ be a $\mathbf{\Gamma}$-isomorphism. Then $f|_X$ is a $\mathbf{\Gamma}$-isomorphism between X and $S = f(X) \in \mathbf{\Gamma}(\mathcal{N})$. □

3.2 Borel and Luzin hierarchies in cb_0-spaces

Here we extend some well known facts on the Borel and Luzin hierarchies in Polish and quasi-Polish spaces.
As is well-known, any uncountable Polish (or quasi-Polish) space is of continuum cardinality. The next fact extends this to many cb_0-spaces:

Proposition 3.5 *Any uncountable space X in $\mathsf{CB}_0(\mathbf{\Sigma}^1_1)$ is of continuum cardinality.*

Proof. By Proposition 3.4, X is $\mathbf{\Sigma}^1_1$-isomorphic to a subspace S of $\mathcal{N}$ such that $S \in \mathbf{\Sigma}^1_1(\mathcal{N})$, so it suffices to show that S is of continuum cardinality. But this follows from a well-known fact of classical DST (Theorem 29.1 in [Ke95]). □

Remark 3.6 The last result can not be improved within ZFC because, as is well known, it is consistent with ZFC that there is a non-countable set $S \in \mathbf{\Pi}^1_1(\mathcal{N})$ of cardinality less than continuum.

Next we establish an extension of the Suslin theorem which equates the Borel sets to the $\mathbf{\Delta}^1_1$-sets. This is a classical result of DST for the case of Polish spaces, and it was extended to quasi-Polish spaces in [Br13].
For this we need the following version of a well-known easy fact:

Lemma 3.7 *Let $X \subseteq Y$ be topological spaces and $1 \leq \alpha < \omega_1$. Then $\mathbf{\Sigma}^1_\alpha(X) = \{X \cap A \mid A \in \mathbf{\Sigma}^1_\alpha(Y)\}$, $\mathbf{\Pi}^1_\alpha(X) = \{X \cap A \mid A \in \mathbf{\Pi}^1_\alpha(Y)\}$, and similarly for the Borel hierarchy.*

Proposition 3.8 *The Suslin theorem holds for any space X in $\mathsf{CB}_0(\mathbf{\Delta}^1_1)$, i.e. $\mathbf{\Delta}^1_1(X) = \bigcup\{\mathbf{\Sigma}^0_\alpha(X) \mid \alpha < \omega_1\}$.*

Proof. It suffices to show the inclusion $\mathbf{\Delta}^1_1(X) \subseteq \bigcup\{\mathbf{\Sigma}^0_\alpha(X) \mid \alpha < \omega_1\}$. Assume without loss of generality that $X \in \mathbf{\Delta}^1_1(P\omega)$. Let $A \in \mathbf{\Delta}^1_1(X)$, then A is in both $\mathbf{\Sigma}^1_1(X)$ and $\mathbf{\Pi}^1_1(X)$. By Lemma 3.7, $A = X \cap B = X \cap C$ for some $B \in \mathbf{\Sigma}^1_1(P\omega)$ and $C \in \mathbf{\Pi}^1_1(P\omega)$. Then $A \in \mathbf{\Delta}^1_1(P\omega)$. By Suslin theorem for $P\omega$, $A \in \mathbf{\Sigma}^0_\alpha(P\omega)$ for some $\alpha < \omega_1$. By Lemma 3.7, $A \in \mathbf{\Sigma}^0_\alpha(X)$. □
As is well known [Ke95], the Borel and Luzin hierarchies do not collapse in any Polish uncountable space X (for the Borel hierarchy, for instance, this means that $\mathbf{\Sigma}^0_\alpha(X) \neq \mathbf{\Pi}^0_\alpha(X)$ for any $\alpha < \omega_1$). In [Br13] this was extended to the quasi-Polish spaces (which coincide with the spaces in $\mathsf{CB}_0(\mathbf{\Pi}^0_2)$). We conclude this section with a further extension of the non-collapse property. For this we need the following lemma:

Lemma 3.9 *Let $1 \leq \alpha < \omega_1$, X, Y be $\mathbf{\Sigma}^0_\alpha$-isomorphic topological spaces, and the Borel hierarchy (resp. the hyperprojective hierarchy) in X does not collapse. Then the Borel hierarchy (resp. the hyperprojective hierarchy) in Y does not collapse.*

Proof. The both hierarchies are treated similarly, so consider only the Borel hierarchy. Suppose for a contradiction that $\mathbf{\Sigma}^0_\beta(Y) = \mathbf{\Pi}^0_\beta(Y)$ for some $\beta < \omega_1$. By the definition of the Borel hierarchy, $\mathbf{\Sigma}^0_\gamma(Y) = \mathbf{\Pi}^0_\gamma(Y)$ for all countable ordinals $\gamma \geq \beta$, in particular for $\gamma = sup\{\alpha, \beta\}$. By Lemma 3.3, $\mathbf{\Sigma}^0_\gamma(X) = \mathbf{\Pi}^0_\gamma(X)$. A contradiction. □

Proposition 3.10 *The Borel and hyperprojective hierarchies do not collapse for any uncountable space X in $\mathsf{CB}_0(\mathbf{\Delta}^1_1)$.*

Proof. By Proposition 3.4 and Lemma 3.9, we can without loss of generality assume that X is a subspace of $\mathcal{N}$ such that $X \in \mathbf{\Delta}^1_1(\mathcal{N})$. By Theorem 29.1 in [Ke95], there is a subspace $C \subseteq X$ homeomorphic to the Cantor space. Since the Borel and hyperprojective hierarchies in C do not collapse, by Lemma 3.7 they also do not collapse in X. □

4 Difference hierarchies

In this section we extend some classical facts on the Hausdorff difference hierarchy (DH) of sets, like the Hausdorff-Kuratowski theorem and the non-collapse property in Polish spaces, to larger classes of cb_0-spaces and to the case of k-partitions.

4.1 Difference hierarchies of sets

Here we recall definition and basic properties of the Hausdorff difference hierarchy of sets, and extend some facts on the DH in Polish spaces to larger classes of cb_0-spaces.
An ordinal α is *even* (resp. *odd*) if $\alpha = \lambda + n$ where λ is either zero or a limit ordinal and $n < \omega$, and the number n is even (resp., odd). For an ordinal α, let $r(\alpha) = 0$ if α is even and $r(\alpha) = 1$, otherwise. For any ordinal α, define the operation D_α sending sequences of sets $\{A_\beta\}_{\beta<\alpha}$ to sets by

$$D_\alpha(\{A_\beta\}_{\beta<\alpha}) = \bigcup\{A_\beta \setminus \bigcup_{\gamma<\beta} A_\gamma \mid \beta < \alpha,\, r(\beta) \neq r(\alpha)\}.$$

For any ordinal $\alpha < \omega_1$ and any pointclass $\mathcal{L}$ in X, let $D_\alpha(\mathcal{L})$ be the class of all sets $D_\alpha(\{A_\beta\}_{\beta<\alpha})$, where $A_\beta \in \mathcal{L}$ for all $\beta < \alpha$. By the *difference hierarchy over* $\mathcal{L}$ we mean the sequence $\{D_\alpha(\mathcal{L})\}_{\alpha<\omega_1}$. Usually we assume that $\mathcal{L}$ is a *base* which by definition means that $\mathcal{L}$ is closed under finite intersection and countable union (note that in finitary versions of the DH we used the term "σ-base" to denote such pointclasses but, since we are interested here only in such pointclasses, we simplify the terminology). As usual, classes $D_\alpha(\mathcal{L})$, $\check{D}_\alpha(\mathcal{L})$ are called non-self-dual levels while $D_\alpha(\mathcal{L}) \cap \check{D}_\alpha(\mathcal{L})$ are called self-dual levels of the DH.
Over bases, the difference hierarchy really looks as a hierarchy, i.e., any level and its dual are contained in all higher levels. The most interesting cases for Descriptive Set Theory are difference hierarchies over non-zero levels of the Borel hierarchy, whose $\mathbf{\Sigma}$-levels are $\mathbf{\Sigma}^{-1,\theta}_\alpha(X) = D_\alpha(\mathbf{\Sigma}^0_\theta(X))$, for any space X and for all $\alpha, \theta < \omega$, $\theta > 0$. For $\theta = 1$, we simplify $\mathbf{\Sigma}^{-1,\theta}_\alpha$ to $\mathbf{\Sigma}^{-1}_\alpha$.
A classical result of DST is the following Hausdorff-Kuratowski theorem:

Theorem 4.1 *Let X be a Polish space. For any non-zero ordinal $\theta < \omega_1$, $\bigcup\{\mathbf{\Sigma}^{-1,\theta}_\alpha(X) \mid \alpha < \omega_1\} = \mathbf{\Delta}^0_{\theta+1}(X)$.*

In [Br13] this result was extended to the quasi-Polish spaces. This extension is an easy corollary of the following nice result (Theorem 68 in [Br13] based on the important Lemma 17 in [SR07]):

Theorem 4.2 *Let X be a cb_0-space, $\delta : D \to X$ an admissible representation of X ($D \subseteq \mathcal{N}$), $A \subseteq X$ X, $\alpha, \theta < \omega_1$ and $\theta \geq 1$. Then $A \in D_\alpha(\mathbf{\Sigma}^0_\theta(X))$ iff $\delta^{-1}(A) \in D_\alpha(\mathbf{\Sigma}^0_\theta(D))$.*

One of the aims of this section is to extend these result from sets to k-partitions. This needs some information on k-forests and h-preorders which are recalled in the next subsection.
Next we establish a partial extension of the Hausdorff-Kuratowski theorem for Polish spaces (which coincide with the spaces in $\mathsf{CB}_0(\mathbf{\Pi}^0_2)$) to a larger class of cb_0-spaces:

Proposition 4.3 *For any space X in $\mathsf{CB}_0(\mathbf{\Delta}^1_1)$ there is a non-zero ordinal $\beta < \omega_1$ such that the Hausdorff-Kuratowski theorem holds in X for each countable ordinal $\theta \geq \beta$, i.e. $\bigcup\{\mathbf{\Sigma}^{-1,\theta}_\alpha(X) \mid \alpha < \omega_1\} = \mathbf{\Delta}^0_{\theta+1}(X)$.*

Proof. Assume without loss of generality that $X \in \mathbf{\Delta}^1_1(P\omega)$. By Suslin theorem for $P\omega$, $X \in \mathbf{\Sigma}^0_\beta(P\omega)$ for some $\beta < \omega_1$. It remains to show that $\mathbf{\Delta}^0_{\theta+1}(X) \subseteq \bigcup\{\mathbf{\Sigma}^{-1,\theta}_\alpha(X) \mid \alpha < \omega_1\}$ for all $\theta \geq \beta$. Let $A \in \mathbf{\Delta}^0_{\theta+1}(X)$, then A is in both $\mathbf{\Sigma}^0_{\theta+1}(X)$ and $\mathbf{\Pi}^0_{\theta+1}(X)$. By Lemma 3.7, $A = X \cap B = X \cap C$ for some $B \in \mathbf{\Sigma}^0_{\theta+1}(P\omega)$ and $C \in \mathbf{\Pi}^0_{\theta+1}(P\omega)$. Then $A \in \mathbf{\Delta}^0_{\theta+1}(P\omega)$. By the Hausdorff-Kuratowski theorem for $P\omega$, $A \in \mathbf{\Sigma}^{-1,\theta}_\alpha(P\omega)$ for some $\alpha < \omega_1$. Since $\theta \geq \beta$ and $X \in \mathbf{\Sigma}^0_\beta(P\omega)$, $A \in \mathbf{\Sigma}^{-1,\theta}_\alpha(X)$. □
The problem of the non-collapse of the DHs is more subtle (compared with the problem of non-collapse of the Borel and Luzin hierarchies) but it is again possible to prove at least a partial result about this property. First we formulate an analogue of Lemma 3.9 which is proved essentially by the same argument:

Lemma 4.4 *Let $1 \leq \alpha < \omega_1$, X, Y be $\mathbf{\Sigma}^0_\alpha$-isomorphic topological spaces, $\alpha \leq \theta < \omega_1$, and the DH $\{\mathbf{\Sigma}^{-1,\theta}_\beta(X)\}_{\beta<\omega_1}$ does not collapse. Then the DH $\{\mathbf{\Sigma}^{-1,\theta}_\beta(Y)\}_{\beta<\omega_1}$ does not collapse.*

Once we have this lemma and note that Lemma 3.7 holds also for the DHs, we easily deduce the following:

Proposition 4.5 *For any uncountable space X in $\mathsf{CB}_0(\mathbf{\Delta}^1_1)$ there is a non-zero ordinal $\alpha < \omega_1$ such that the DH $\{\mathbf{\Sigma}^{-1,\theta}_\beta(X)\}_{\beta<\omega_1}$ does not collapse for each countable ordinal $\theta \geq \alpha$.*

4.2 h-Preorder

Here we discuss some posets which serve as notation systems for levels of the DHs of k-partitions. All notions and facts of this subsection are contained (at least, implicitly) in [Se07, Se07a].
Posets considered here are assumed to be (at most) countable and without infinite chains. The absence of infinite chains in a poset $(P;\leq)$ is of course equivalent to well-foundednes of both $(P;\leq)$ and $(P;\geq)$. By a *forest* we mean a poset without infinite chains in which every upper cone $\{y \mid x \leq y\}$ is a chain. *Tree* is a forest having the biggest element (called *the root* of the tree).
Let $(T;\leq)$ be a tree without infinite chains; in particular, it is well-founded. As for each well-founded partial order, there is a canonical rank function rk_T from T to ordinals. The rank $rk(T)$ of $(T;\leq)$ is by definition the ordinal $rk_T(r)$, where r is the root of $(T;\leq)$. It is well-known that the rank of any countable tree without infinite chains is a countable ordinal, and any countable ordinal is the rank of such a tree.
A *k-poset* is a triple $(P;\leq,c)$ consisting of a poset $(P;\leq)$ and a labeling $c : P \to k$. Rank of a k-poset $(T;\leq,c)$ is by definition the rank of $(T;\leq)$. *Morphism* $f : (P;\leq,c) \to (P';\leq',c')$ between k-posets is a monotone function $f : (P;\leq) \to (P';\leq')$ respecting the labelings, i.e. satisfying $c = c' \circ f$. Let $\widetilde{\mathcal{F}}_k$ and $\widetilde{\mathcal{T}}_k$ denote the classes of all countable k-forests and countable k-trees without infinite chains, respectively. Note that we use tilde in our notation in order to distinguish the introduced objects from their finitary versions extensively studied in [Se06, Se08a, Se12].
The h-preorder $\leq_h$ on $\widetilde{\mathcal{P}}_k$ is defined as follows: $(P,\leq,c) \leq_h (P',\leq',c')$, if there is a morphism from $(P,\leq,c)$ to $(P',\leq',c')$. Let $\widetilde{\mathbb{F}}_k, \widetilde{\mathbb{T}}_k$ be the quotient posets of $\widetilde{\mathcal{F}}_k$ and $\widetilde{\mathcal{T}}_k$ under the h-equivalence, respectively. Let $\widetilde{\mathbb{F}}'_k$ be obtained from $\widetilde{\mathbb{F}}_k$ by adjoining a new smallest element $\perp$ (corresponding to the empty forest). Let $P \sqcup Q$ be the disjoint union of k-posets P, Q and $\bigsqcup_i P_i = P_0 \sqcup P_1 \sqcup \cdots$ the disjoint union of an infinite sequence $P_0, P_1, \ldots$ of k-posets. For a k-forest F and $i < k$, let $p_i(F)$ be the k-tree obtained from F by adjoining a new biggest element and assigning the label i to this element. It is clear that the introduced operations respect the h-equivalence and that any countable k-forest is h-equivalent to a countable term of signature $\{\sqcup, p_0, \ldots, p_{k-1}, 0, \ldots, k-1\}$ without free variables (the constant symbol i in the signature is interpreted as the singleton tree carrying the label i).

Proposition 4.6 (1) *For any $k \geq 2$, the structures $(\widetilde{\mathcal{F}}_k;\leq)$ and $(\widetilde{\mathcal{T}}_k;\leq)$ are well preorders of rank ω_1.*

(2) *The posets $\widetilde{\mathbb{F}}_2$ and $\widetilde{\mathbb{T}}_2$ have width 2 (i.e., they have no antichains with more that 2 elements).*

(3) *The poset $\widetilde{\mathbb{F}}'_k$ is a distributive lattice where any countable set of elements have a supremum.*

(4) *The set $\sigma ji(\widetilde{\mathbb{F}}'_k)$ of σ-join-irreducible elements of the lattice $\widetilde{\mathbb{F}}'_k$ (i.e., the elements x such that $x \leq \bigsqcup\{y_n \mid n < \omega\}$ implies that $y \leq y_n$ for some n) coincides with $\widetilde{\mathbb{T}}_k$.*

(5) *The set $ji(\widetilde{\mathbb{F}}'_k)$ of join-irreducible elements of the lattice $\widetilde{\mathbb{F}}'_k$ coincides with $\widetilde{\mathbb{T}}_k \cup S$ where S is the set of supremums of infinite increasing sequences of elements in $\widetilde{\mathbb{T}}_k$.*

(6) *Any element of $\widetilde{\mathbb{F}}'_k$ is the infimum of finitely many elements of $\widetilde{\mathbb{T}}_k$.*

For a result in the next subsection we need the following canonical representatives for the structures $\widetilde{\mathcal{F}}_2$ and $\widetilde{\mathcal{T}}_2$. Define by induction the sequence $\{T_\alpha\}_{\alpha<\omega_1}$ as follows: $T_0 = 0$, $T_{\alpha+1} = p_0(\overline{T}_\alpha)$ where $\overline{T}_\alpha$ is obtained from T_α by changing any label $l < 2$ by the label $1-l$, and $T_\lambda = p_0(\overline{T}_{\alpha_0} \sqcup \overline{T}_{\alpha_1} \sqcup \cdots)$ for a limit ordinal λ where $\alpha_0 < \alpha_1 < \cdots$ is a sequence of odd ordinals satisfying $sup\{\alpha_n \mid n < \omega\} = \lambda$. The next assertion follows from the proof of the corresponding result in [Se07, Se07a].

Proposition 4.7 (1) *For all $\alpha < \beta < \omega_1$, $T_\alpha \sqcup \overline{T}_\alpha <_h T_\beta$ and $T_\alpha, \overline{T}_\alpha$ are $\leq_h$-incomparable.*

(2) *Any element of $\widetilde{\mathcal{T}}_2$ (resp. of $\widetilde{\mathcal{F}}_2$) is h-equivalent to precisely one of $T_\alpha, \overline{T}_\alpha$, (resp. to precisely one of $T_\alpha, \overline{T}_\alpha, T_\alpha \sqcup \overline{T}_\alpha, \bigsqcup_{\alpha<\lambda} T_\alpha$ where λ is a limit countable ordinal).*

(3) *$T_\alpha \sqcup \overline{T}_\alpha \equiv_h T_{\alpha+1} \sqcap \overline{T}_{\alpha+1}$ and $\bigsqcup_{\alpha<\lambda} T_\alpha \equiv_h T_\lambda \sqcap \overline{T}_\lambda$ for each limit countable ordinal λ, where $\sqcap$ is the binary operation of k-forests inducing the infimum operation in $\widetilde{\mathbb{F}}'_k$.*

4.3 Difference hierarchies of k-partitions

Here we extend the difference hierarchy of sets to that of k-partitions. Note that similar hierarchies were considered e.g. in [Se07, Se07a] but in fact the definition here is slightly different which results in equivalence of the corresponding DHs only over the bases with ω-reduction properties. In general, the definition here is better than the previous one because it yields, for example, Theorem 4.14 for $\theta = 1$ which is not always the case for the previous definition.

Let $\mathcal{L}$ be a base in X. Recall that a k-partition of X is a function $A : X \to k$ often written as a tuple $(A_0, \dots, A_{k-1})$ where $A_i = \{x \in X \mid A(x) = i\}$. By a *partial k-partition of X* we mean a function $A : Y \to k$ for some $Y \in \mathcal{L}$. Let $P \in \widetilde{\mathcal{F}}_k$. We say that a partial k-partition A is defined by a P-family $\{B_p\}_{p\in P}$ of $\mathcal{L}$-sets if $A_i = \bigcup_{p\in P_i} \tilde{B}_p$ for each $i < k$ where $\tilde{B}_p = B_p \setminus \bigcup_{q<p} B_q$ and $P_i = c^{-1}(i)$; note that in this case $A \in k^Y$ where $Y = \bigcup_{p\in P} B_p$.

We denote by $\mathcal{L}^Y(P)$ the set of partitions $A : Y \to k$ defined by some P-family $\{B_p\}_{p\in P}$ of $\mathcal{L}$-sets. In case $Y = X$ we omit the superscript X and call the classes $\mathcal{L}(P), P \in \widetilde{\mathcal{F}}_k$ *levels of the DH of k-partitions over $\mathcal{L}$*. We formulate some basic properties of the levels. We omit the proofs because they are quite similar to the corresponding proofs for the finitary version of the DH in [Se12].

Proposition 4.8 (1) *If $A \in \mathcal{L}^Y(P)$ then $A|_Z \in \mathcal{L}^Z(P)$ for each $Z \subseteq Y$, $Z \in \mathcal{L}$.*

(2) *Any $A \in \mathcal{L}^Y(P)$ is defined by a monotone P-family $\{C_p\}$ (monotonicity means that $C_q \subseteq C_p$ for $q \le p$).*

(3) *Let f be a function on X such that $f^{-1}(A) \in \mathcal{L}$ for each $A \in \mathcal{L}$. Then $A \in \mathcal{L}^Y(P)$ implies $f^{-1}(A) = (f^{-1}(A_0), \dots, f^{-1}(A_{k-1})) \in \mathcal{L}^{f^{-1}(Y)}(P)$.*

(4) *If $P \le_h Q$ then $\mathcal{L}^Y(P) \subseteq \mathcal{L}^Y(Q)$.*

(5) *For all $F, G \in \widetilde{\mathcal{F}}_k$, $\mathcal{L}(F) \cap \mathcal{L}(G) = \mathcal{L}(F \sqcap G)$.*

Proposition 4.8(5) and Proposition 4.6(6) show that any level $\mathcal{L}(F)$ is a finite intersection of the levels $\mathcal{L}(T)$, $T \in \widetilde{\mathcal{T}}_k$. This remark, together with the results below, suggest that the levels $\mathcal{L}(T)$, $T \in \widetilde{\mathcal{T}}_k$ are analogs for the DH of k-partitions of the non-self-dual levels $D_\alpha(\mathcal{L}), \check{D}_\alpha(\mathcal{L})$ of the DH of sets. Therefore, the precise analog of the DH of sets is the family $\{\mathcal{L}(T)\}_{T\in\widetilde{\mathcal{T}}_k}$ rather than the family $\{\mathcal{L}(F)\}_{F\in\widetilde{\mathcal{F}}_k}$.

The meaning of the last paragraph might be not clear because it is not even obvious that the DH of k-partitions really extends the DH of sets; we have at least to show that the DH of 2-partitions essentially coincides with the DH of sets. We do this in the next proposition where we employ the 2-trees T_α from Subsection 4.2.

Proposition 4.9 *Let $\mathcal{L}$ be a base in X. Then $\mathcal{L}(T_\alpha) = D_\alpha(\mathcal{L})$ for each $\alpha < \omega_1$.*

Proof. Let $A \in \mathcal{L}(T_\alpha)$ be defined by a family $\{B_p\}_{p\in T_\alpha}$ where $B_p \in \mathcal{L}$. Define the sequence $\{A_\beta\}_{\beta<\alpha}$ as follows: if $r(\beta) = r(\alpha)$ then $A_\beta = \bigcup\{B_p \mid rk(p) \le \beta \wedge c(p) = 0\}$, otherwise $A_\beta = \bigcup\{B_p \mid rk(p) \le \beta \wedge c(p) = 1\}$. Then $A_\beta \in \mathcal{L}$ and $A = \{\tilde{A}_\beta \mid r(\beta) \ne r(\alpha)\}$ where $\tilde{A}_\beta = A_\beta \setminus \bigcup_{\gamma<\beta} A_\gamma$, hence $A \in D_\alpha(\mathcal{L})$.

Conversely, let $A \in D_\alpha(\mathcal{L})$, then $A = \{\tilde{A}_\beta \mid r(\beta) \ne r(\alpha)\}$ for some sequence $\{A_\beta\}_{\beta<\alpha}$ of $\mathcal{L}$-sets. Define the family $\{B_p\}_{p\in T_\alpha}$ of $\mathcal{L}$-sets as follows: if p is the root of T_α then $B_p = X$, otherwise $B_p = A_{rk(p)}$ where $rk : T_\alpha \to \alpha + 1$ is the rank function. Since rk is a surjection for each α, $\tilde{B}_p = \tilde{A}_{rk(p)}$.

Note that $\tilde{B}_p, \tilde{B}_q$ are disjoint whenever $c(p) \ne c(q)$ because $\tilde{A}_\beta, \tilde{A}_\gamma$ are disjoint for distinct $\beta, \gamma < \alpha$. If p is the root of T_α then $\tilde{B}_p = X \setminus \bigcup_{\beta<\alpha} \tilde{A}_\beta \subseteq \overline{A}$. If p is not the root then, by the definition of T_α, $r(rk(p)) = r(\alpha)$ iff $c(p) = 0$. Then for each $x \in \bigcup_{\beta<\alpha} \tilde{A}_\beta$ we have: $x \in A$ iff $r(\beta) \ne r(\alpha)$ (where β is the unique ordinal with $x \in \tilde{A}_\beta$) iff $c(p) = 1$. Thus, $A \in \mathcal{L}(T_\alpha)$ is defined by $\{B_p\}_{p\in T_\alpha}$ and therefore $A \in \mathcal{L}(T_\alpha)$. □

For $F \in \widetilde{\mathcal{F}}_k$, by a *reduced F-family of $\mathcal{L}$-sets* we mean a monotone F-family $\{B_p\}_{p\in F}$ of $\mathcal{L}$-sets such that $B_p \cap B_q = \emptyset$ for all incomparable $p, q \in F$. Let $A \in \mathcal{L}^Y_r(F)$ be the set of partial k-partitions defined by reduced F-families $\{B_p\}_{p\in F}$ of $\mathcal{L}$-sets such that $\bigcup_p B_p = Y$. The next result is an infinitary version of Proposition 7.15 [Se12] and is proved by essentially the same argument.

Proposition 4.10 *Let $\mathcal{L}$ have the ω-reduction property, $Y \in \mathcal{L}$ and $F \in \mathcal{F}_k$. Then $\mathcal{L}^Y(F) = \mathcal{L}^Y_r(F)$, in particular $\mathcal{L}(F) = \mathcal{L}_r(F)$.*

In [Se13], principal total representations (TR) of pointclasses were introduced and studied. The results in [Se13] naturally extend to k-partitions. By a *family of k-partition classes* we mean a family $\mathbf{\Gamma} = \{\mathbf{\Gamma}(X)\}$ indexed by arbitrary topological spaces X such that $\mathbf{\Gamma}(X) \subseteq k^X$ and $A \circ f \in \mathbf{\Gamma}(X)$ for any continuous function $f : X \to Y$ and any k-partition $A : Y \to k$ from $\mathbf{\Gamma}(Y)$.
We note that, by Proposition 4.8(3), the levels of the DHs over any $\mathbf{\Sigma}$-level of the Borel or Luzin hierarchy are families of k-partition classes. Let $\mathbf{\Sigma}^0_\theta(X, F)$ be the F-level ($F \in \widetilde{\mathcal{F}}_k$) of the DH of k-partitions over $\mathbf{\Sigma}^0_\theta(X)$. The next fact is obvious:

Proposition 4.11 *Let θ be a non-zero countable ordinal and $F \in \widetilde{\mathcal{F}}_k$. Then $\mathbf{\Sigma}^0_\theta(F) = \{\mathbf{\Sigma}^0_\theta(X, F)\}_X$ is a family of k-partition classes.*

Let $\{\mathbf{\Gamma}(X)\}$ be a family of k-partition classes. A function $\nu : \mathcal{N} \to \mathbf{\Gamma}(X)$ is a *$\mathbf{\Gamma}$-TR* if the k-partition $\lambda a, x.\nu(a)(x)$ is in $\mathbf{\Gamma}(\mathcal{N} \times X)$. Such ν is a *principal $\mathbf{\Gamma}$-TR* if any $\mathbf{\Gamma}$-TR $\mu : \mathcal{N} \to \mathbf{\Gamma}(X)$ is continuously reducible to ν.
According to Theorem 5.2 in [Se13], the non-self-dual levels $\mathbf{\Gamma}$-TR of the classical hierarchies in cb_0-spaces have $\mathbf{\Gamma}(X)$-TRs. This result extends to the "non-self-dual" levels of the DHs of k-partitions but only under the additional assumption that the corresponding bases have the ω-reduction property:

Proposition 4.12 *Let X be a cb_0-space, $\theta \geq 2$ a countable ordinal and $T \in \widetilde{\mathcal{T}}_k$. Then $\mathbf{\Sigma}^0_\theta(X, T)$ has a principal $\mathbf{\Sigma}^0_\theta(T)$-TR. If X is in addition zero-dimensional then $\mathbf{\Sigma}^0_1(X, T)$ has a principal $\mathbf{\Sigma}^0_1(T)$-TR.*

Proof Hint. Modulo Theorem 5.2 in [Se13], the proof is straightforward, so we give only informal proof hints. Note that, by Proposition 2.3, the assumptions on θ guarantee that the class $\mathbf{\Sigma}^0_\theta(X)$ has the ω-reduction property, so by Proposition 4.10 the elements of $\mathbf{\Sigma}^0_\theta(X, T)$ are precisely those defined by the monotone reduced T-families $\{B_p\}$ of $\mathbf{\Sigma}^0_\theta(X)$-sets with $B_p = X$ where p is the root of T. The principal $\mathbf{\Sigma}^0_\theta$-TR of $\mathbf{\Sigma}^0_\theta(X)$ induces a natural representaion of all T-families $\{C_p\}$ of $\mathbf{\Sigma}^0_\theta(X)$-sets with $C_p = X$. The problem is that such a family typically does not define any k-partition, hence we do not in general have an induced TR of $\mathbf{\Sigma}^0_\theta(X, T)$. But the ω-reduction property gives a uniform procedure of transforming $\{C_p\}$ to a monotone reduced T-family $\{B_p\}$ of $\mathbf{\Sigma}^0_\theta(X)$-sets with $B_p = X$, such that $\{B_p\} = \{C_p\}$ whenever $\{C_p\}$ already has this property. Since any such $\{B_p\}$ defines an element of $\mathbf{\Sigma}^0_\theta(X, T)$, this induces a TR of $\mathbf{\Sigma}^0_\theta(X, T)$. It is straightforward to check that this TR has the desired properties. □
We conclude this section with extending the Hausdorff-Kuratowski theorem to k-partitions. First we extend Theorem 4.2 to k-partitions:

Theorem 4.13 *Let X be a cb_0-space, $\delta : D \to X$ an admissible representation of X ($D \subseteq \mathcal{N}$), $A : X \to k$ a k-partition of X, $\theta \geq 1$ a countable ordinal and $F \in \widetilde{\mathcal{F}}_k$. Then $A \in \mathbf{\Sigma}^0_\theta(X, F)$ iff $A \circ \delta \in \mathbf{\Sigma}^0_\theta(D, F)$.*

Proof is similar to the proof of Theorem 68 in [Br13]. Let first $A \in \mathbf{\Sigma}^0_\theta(X, T)$, then A is defined by an F-family $\{B_p\}$ of $\mathbf{\Sigma}^0_\theta(X)$-sets. Then $A \circ \delta$ is defined by the F-family $\{\delta^{-1}(B_p)\}$ of $\mathbf{\Sigma}^0_\theta(D)$-sets, hence $A \circ \delta \in \mathbf{\Sigma}^0_\theta(D, T)$.
Conversely, let $A \circ \delta \in \mathbf{\Sigma}^0_\theta(D, T)$, then $A \circ \delta$ is defined by an F-family $\{C_p\}$ of $\mathbf{\Sigma}^0_\theta(D)$-sets, so $\delta^{-1}(A_i) = \bigcup\{\tilde{C}_p \mid c(p) = i\}$ for each $i < k$. By the proof of Theorem 68 in [Br13], we can without loss of generality assume that δ has Polish fibers, i.e., $\delta^{-1}(x)$ is Polish for each $x \in X$. For any $p \in F$, let B_p consist of the elements $x \in X$ such that the set $C_p \cap \delta^{-1}(x)$ is non-meager in $\delta^{-1}(x)$. By the proof of Theorem 68 in [Br13], $B_p \in \mathbf{\Sigma}^0_\theta(X)$, hence it suffices to show that A is defined by the F-family $\{B_p\}$.
First we check that $\tilde{B}_p \subseteq \delta(\tilde{C}_p)$. Let $x \in \tilde{B}_p$, so $C_p \cap \delta^{-1}(x)$ is non-meager in $\delta^{-1}(x)$ and $C_q \cap \delta^{-1}(x)$ is meager in $\delta^{-1}(x)$ for each $q < p$, hence also $(\bigcup_{q<p} C_q) \cap \delta^{-1}(x)$ is meager in $\delta^{-1}(x)$. Since

$$C_p \cap \delta^{-1}(x) = (\tilde{C}_p \cap \delta^{-1}(x)) \cup (\bigcup_{q<p} C_q) \cap \delta^{-1}(x),$$

$\tilde{C}_p \cap \delta^{-1}(x)$ is non-meager in $\delta^{-1}(x)$, in particular $\tilde{C}_p \cap \delta^{-1}(x)$ is non-empty. Let $a \in \tilde{C}_p \cap \delta^{-1}(x)$, then $x = \delta(a) \in \delta(\tilde{C}_p)$, as desired.
We have to show that $A_i = \bigcup\{\tilde{B}_p \mid c(p) = i\}$ for each $i < k$. Let first $x \in \tilde{B}_p$, $c(p) = i$. Then $x = \delta(a)$ for some $a \in \tilde{C}_p \subseteq \delta^{-1}(A_i)$. Thus, $x \in A_i$.

Conversely, let $x \in A_i$. Choose $a \in D$ with $x = \delta(a)$. Then $a \in \delta^{-1}(A_i)$, so $a \in \tilde{C}_p$ for some $p \in F$, $c(p) = i$. Note that $x \in B_q$ for some $q \in F$ (otherwise, $C_q \cap \delta^{-1}(x)$ is meager in $\delta^{-1}(x)$ for each $q \in F$, hence also $\delta^{-1}(x)$ is meager, a contradiction). Then $x \in \tilde{B}_q$ for some $q \in F$, hence $x = \delta(b)$ for some $b \in \tilde{C}_q$. Let $j = c(q)$, then $a \in \delta^{-1}(A_i)$ and $b \in \delta^{-1}(A_j)$, then $x \in A_i \cap A_j$, so $i = j$ and $c(q) = i$. $\square$
As an immediate corollary, we obtain the Hausdorff-Kuratowski theorem for k-partitions in quasi-Polish spaces.

Theorem 4.14 *Let X be a quasi-Polish space and $\theta \geq 1$ a countable ordinal. Then $\bigcup\{\mathbf{\Sigma}^0_\theta(X,T) \mid F \in \widetilde{\mathcal{F}}_k\} = (\mathbf{\Delta}^0_{\theta+1}(X))_k$.*

Proof. For $\theta \geq 2$, the assertion follows from Theorem 5.1 in [Se08] and Theorem 4.2 but for $\theta = 1$ the result is new. Let $A \in (\mathbf{\Delta}^0_2(X))_k$, we have to show that $A \in \bigcup\{\mathbf{\Sigma}^0_1(X,T) \mid F \in \widetilde{\mathcal{F}}_k\}$. Let δ be an admissible total representation of X. Then $A \circ \delta \in (\mathbf{\Delta}^0_2(\mathcal{N}))_k$. Since $\mathbf{\Sigma}^0_1(\mathcal{N})$ has the ω-reduction property, $A \circ \delta \in \bigcup\{\mathbf{\Sigma}^0_\alpha(\mathcal{N},F) \mid F \in \widetilde{\mathcal{F}}_k\}$ by Theorem 5.1 in [Se08]. By Theorem 4.13, $A \in \bigcup\{\mathbf{\Sigma}^0_1(X,T) \mid F \in \widetilde{\mathcal{F}}_k\}$. $\square$

Remark 4.15 Using the argument of Proposition 4.3, it is straightforward to extend the last result to spaces X in $\mathsf{CB}_0(\mathbf{\Delta}^1_1)$.

5 Fine hierarchies

In this section we extend the DH of k-partitions to the FH of k-partitions. Many results and proofs here extend the ones for the DH from the previous section or the corresponding results on the finitary version of the FH from [Se12], so we concentrate only on the new moments and try to be concise whenever the material is a straightforward extension of the previous one. We will see that the FH of k-partitions is in a sense an "iterated version" of the DH of k-partitions which is far from obvious for the the particular case of the Wadge hierarchy, under the classical definition.
As we explain below, the FH of sets in the Baire space does coincide with the Wadge hierarchy. To our knowledge, the extension of this hierarchy to non-zero-dimensional spaces is new here (so far such an extension was known only for the finitatry version of the FH [Se08a, Se12]). Interestingly, in our approach here the definition of the FH of sets is in fact not simpler than for the k-partitions.
Since even the definition of the FH is technically very involved, we concentrate here on a slightly easier case of sets and k-partitions of finite Borel rank and provide only proof hints for some long proofs, appealing to the analogy with the finitary version in [Se12].

5.1 More on the h-preorders

Here we extend some notions and facts from Subsection 4.2 about the h-preorders, in order to describe notation systems for levels of the FHs of k-partitions. We omit the proofs which are easy variations of their finitary versions in [Se12].
Let $(Q;\leq)$ be a poset. A *Q-poset* is a triple $(P,\leq,c)$ consisting of a countable nonempty poset $(P;\leq)$, $P \subseteq \omega$ without infinite chains, and a labeling $c : P \to Q$. By default, we denote the labeling in a Q-poset by c. A *morphism* $f : (P,\leq,c) \to (P',\leq',c')$ of Q-posets is a monotone function $f : (P;\leq) \to (P';\leq')$ satisfying $\forall x \in P(c(x)) \leq c'(f(x)))$. Let $\widetilde{\mathcal{P}}_Q$, $\widetilde{\mathcal{F}}_Q$ and $\widetilde{\mathcal{T}}_Q$ denote the sets of all countable Q-posets, Q-forests and Q-trees, respectively.
The *h-preorder* $\leq_h$ on $\mathcal{P}_Q$ is defined as follows: $P \leq_h P'$, if there is a morphism from P to P'. The quotient-posets of $\widetilde{\mathcal{P}}_Q$, $\widetilde{\mathcal{F}}_Q$, $\widetilde{\mathcal{T}}_Q$ are denoted $\widetilde{\mathbb{P}}_Q$, $\widetilde{\mathbb{F}}_Q$, $\widetilde{\mathbb{T}}_Q$, respectively. Note that for the particular case $Q = \bar{k}$ of the antichain with k elements we obtain the preorders $\widetilde{\mathcal{P}}_k$, $\widetilde{\mathcal{F}}_k$, $\widetilde{\mathcal{T}}_k$ from the previous section.
Next we formulate some lattice-theoretic properties of the h-preorders. By a *partial lower semilattice* we mean a poset in which any two elements that have a lower bound have a (unique) greatest lower bound. For any poset Q, let Q' be the poset obtained from Q by adjoining the new element $\perp$ which is smaller than all elements in Q. Define the function $s : Q \to \mathcal{P}_Q$ as follows: $s(q)$ is the singleton tree labeled by $q \in Q$. If Q is an upper semilattice, define the function $l : \mathcal{P}_Q \to Q$ by $l(P) = \bigcup\{c(p) \mid p \in P\}$ where $\cup$

is the join operation in Q. For Q-posets P and R (resp. $P_0, P_1, \ldots$), let $P \sqcup R$ (resp. $\bigcup_i P_i$) denote their disjoint union (resp. countable disjoint union), $P \sqcup R, \bigcup_i P_i \in \widetilde{\mathcal{P}}_Q$.
For posets P and Q we write $P \subseteq Q$ (resp. $P \sqsubseteq Q$) if P is a substructure of Q (resp. P is an initial segment of Q). The next assertion is essentially contained in [Se12].

Proposition 5.1 (1) *If Q is a partial lower semilattice then Q' is a lower semilattice.*

(2) *$(\widetilde{\mathcal{P}}_Q; \leq_h, \sqcup)$ is a distributive upper σ-semilattice that contains $(\mathcal{F}_Q; \leq_h, \sqcup)$ as a distributive upper σ-subsemilattice.*

(3) *If Q is a partial lower semilattice then $(\widetilde{\mathcal{P}}'_Q; \leq_h)$ and $(\widetilde{\mathcal{F}}'_Q; \leq_h)$ are distributive lattices.*

(4) *If Q is directed (i.e. any two elements have an upper bound) then $s(Q)$ is a cofinal subset of $\widetilde{\mathcal{P}}_Q$ (i.e. any $x \in \widetilde{\mathcal{P}}_Q$ is below $s(q)$ for some $q \in Q$).*

(5) *The mapping s is an isomorphic embedding of Q into $\widetilde{\mathcal{P}}_Q$.*

(6) *If Q is a partial lower semilattice then s preserves the greatest lower bound operations in Q and $\widetilde{\mathcal{P}}_Q$ (and similarly for $\widetilde{\mathcal{F}}_Q$).*

(7) *If Q is an upper semilattice then $l : \widetilde{\mathcal{P}}_Q \to Q$ is a homomorphism of upper semilattices and $q = l(s(q))$ for each $q \in Q$.*

(8) *If Q is a bqo then $(\widetilde{\mathcal{F}}_Q; \leq_h)$, $(\widetilde{\mathcal{T}}_Q; \leq_h)$ are bqo's.*

(9) *Let P, Q be arbitrary posets. Then $P \subseteq Q$ implies $\widetilde{\mathbb{F}}_P \subseteq \widetilde{\mathbb{F}}_Q$, and $P \sqsubseteq Q$ implies $\widetilde{\mathbb{F}}_P \sqsubseteq \widetilde{\mathbb{F}}_Q$.*

Now we can iterate the construction $Q \mapsto \widetilde{\mathbb{F}}_Q$ starting with the antichain $\overline{k}$ of k elements $\{0, \ldots, k-1\}$. Define the sequence $\{\widetilde{\mathcal{F}}_k(n)\}_{n<\omega}$ of preorders by induction on n as follows: $\widetilde{\mathcal{F}}_k(0) = \overline{k}$ and $\widetilde{\mathcal{F}}_k(n+1) = \widetilde{\mathcal{F}}_{\widetilde{\mathcal{F}}_k(n)}$. Identifying the elements $i < k$ of $\overline{k}$ with the corresponding minimal elements $s(i)$ of $\widetilde{\mathcal{F}}_k(1)$, we may think that $\widetilde{\mathcal{F}}_k(0) \sqsubseteq \widetilde{\mathcal{F}}_k(1)$. By items (8,9) of Proposition 5.1, $\widetilde{\mathcal{F}}_k(n) \sqsubseteq \widetilde{\mathcal{F}}_k(n+1)$ for each $n < \omega$, and $\widetilde{\mathcal{F}}_k(\omega) = \bigcup_{n<\omega} \widetilde{\mathcal{F}}_k(n)$ is a wqo. For any $n \leq \omega$, let $\widetilde{\mathbb{F}}_k(n)$ be the quotient-poset of $\widetilde{\mathcal{F}}_k(n)$.
Of course, similar constructions can be done with $\widetilde{\mathcal{T}}$ in place of $\widetilde{\mathcal{F}}$. The preorders $\widetilde{\mathcal{F}}_k(\omega)$, $\widetilde{\mathcal{T}}_k(\omega)$ and the set $\widetilde{\mathcal{T}}^{\sqcup}_k(\omega)$ of countable joins of elements in $\widetilde{\mathcal{T}}_k(\omega)$, play an important role in the study of the FH of k-partitions because they provide convenient naming systems for the levels of this hierarchy (similar to the previous section where $\widetilde{\mathcal{F}}_k$ and $\widetilde{\mathcal{T}}_k$ where used to name the levels of the DH of k-partitions). Note that $\widetilde{\mathcal{F}}_k(1) = \widetilde{\mathcal{F}}_k$ and $\widetilde{\mathcal{T}}_k(1) = \widetilde{\mathcal{T}}_k$.
By Proposition 5.1, for any $n < \omega$ there is an embedding $s = s_n$ of $\widetilde{\mathbb{F}}_k(n)$ into $\widetilde{\mathbb{F}}_k(n+1)$, and s_{n+1} coincides with s_n on $\widetilde{\mathbb{F}}_k(n)$. This induces the embedding $s = \bigcup_{n<\omega} s_n$ of $\mathbb{F}_k(\omega)$ into itself such that s coincides with s_n on $\widetilde{\mathbb{F}}_k(n)$ for each $n < \omega$. Similarly, for any $n < \omega$ we have the function l from $\widetilde{\mathbb{F}}_k(n+2)$ onto $\widetilde{\mathbb{F}}_k(n+1)$ which induces the function (denoted also by l) from $\widetilde{\mathbb{F}}_k(\omega)$ onto $\widetilde{\mathbb{F}}_k(\omega)$.
Define the binary operation $*$ on $\widetilde{\mathcal{P}}_k(\omega)$ as follows: $F * G$ is the labeled poset obtained from G by adjoining a new largest (root) element and assigning the label F to that element. It is easy to see that the operation $*$ respects the h-equivalence relation and hence induces the binary operation on $\widetilde{\mathbb{P}}_k(\omega)$ also denoted by $*$. Note that for $F = s(i) = i < k$ we have $F * G = p_i(G)$ and that $\widetilde{\mathcal{F}}_k(\omega)$ is closed under $*$.
We formulate some properties of the introduced objects illustrating a rich algebraic structure of $\widetilde{\mathcal{F}}_k(\omega)$. They are again almost the same as their corresponding finitary versions in [Se12].

Proposition 5.2 (1) *For any n with $1 \leq n \leq \omega$, $\widetilde{\mathbb{F}}'_k(n)$ is a well distributive lattice.*

(2) *Any element of $\widetilde{\mathbb{F}}'_k(\omega)$ is the value of a variable-free term (countable joins are allowed) of signature $\{\sqcup, *, \bot, 0, \ldots, k-1\}$.*

(3) *For any $0 < n \leq \omega$, $\widetilde{\mathbb{T}}_k(n)$ generates $\widetilde{\mathbb{F}}'_k(n)$ under $\sqcap$.*

(4) *The set of σ-join-irreducible elements of $\widetilde{\mathbb{T}}^{\sqcup}_k(\omega)$ coincides with $\widetilde{\mathbb{T}}_k(\omega)$.*

(5) *The set of join-irreducible elements $\mathbb{T}^{\sqcup}_k(\omega)$ coincides with $\widetilde{\mathbb{T}}_k(\omega) \cup S$, where S is the set of supremums of infinite increasing sequences in $\widetilde{\mathbb{T}}_k(\omega)$.*

Next we provide a characterization of $\widetilde{\mathbb{P}}_k(\omega)$ (and the related substructures) which is sometimes more convenient when dealing with the FH in the next subsection. For $P, Q \in \widetilde{\mathcal{P}}_k(n)$, an *explicit morphism*

$\varphi : P \to Q$ is a sequence $(\varphi_0, \ldots, \varphi_{n-1})$ where φ_0 is a morphism from P to Q, $\varphi_1 = \{\varphi_{1,p_0}\}_{p_0 \in P}$ is a family of morphisms from $c(p_0)$ to $c(\varphi_0(p_0))$, $\varphi_2 = \{\varphi_{2,p_0,p_1}\}_{p_0 \in P, p_1 \in c(p_0)}$ is a family of morphisms from $c(p_1)$ to $c(\varphi_{1,p_0}(p_1))$ and so on (this notion makes use of the convention that $i = s(i)$ for each $i < k$). Note that $P \leq_h Q$ iff there is an explicit morphism from P ro Q, and that for $n = 1$ the explicit morphisms essentially coincide with the morphisms. In the next assertion we treat $\widetilde{\mathbb{P}}_k(\omega)$ as the category whose morphisms are the explicit morphisms.

For each positive $n < \omega$, a *k-labeled n-preorder* [Se08a] is a countable structure $(S; d, \leq_0, \cdots, \leq_{n-1})$ where $d : S \to k$ is a k-partition of S and $\leq_0, \cdots, \leq_n$ are preorders on S such that $\leq_n$ is a partial order, $x \leq_{i+1} y$ implies $x \equiv_i y$, the quotient-poset of $(S; \leq_0)$ has no infinite chains, for each $x \in S$ the quotient-poset of $([x]_0; \leq_1)$ has no infinite chains for each $y \in [x]_0$ the quotient-poset of $([y]_1; \leq_2)$ has no infinite chains, and so on. Let $\mathcal{S}_n(k)$ be the category of k-labeled n-posets as objects where the morphisms are functions that preserve the labelings and are monotone w.r.t. all the preorders.

Proposition 5.3 *For any positive $n < \omega$, the categories $\widetilde{\mathcal{P}}_k(n)$ and $\mathcal{S}_n(k)$ are equivalent.*

Proof Hint. The equivalence is witnessed (cf. Proposition 8.8 of [Se12]) by the functors defined as follows. Relate to any object $(P; \leq, c)$ of $\widetilde{\mathcal{P}}_k(n)$ the object $P^\circ = (X; \leq_0, \ldots, \leq_{n-1}, d)$ of $\mathcal{S}_n(k)$ where X is formed by the elements $p = (p_0, \ldots, p_{n-1})$ such that $p_0 \in P, p_1 \in c(p_0), \ldots$, the preorders between such p and $r = (r_0, \ldots, r_{n-1})$ are defined by $p \leq_0 r$ iff $p_0 \leq r_0$, $p \leq_1 r$ iff $p_0 = r_0$ and $p_1 \leq r_1$ and so on, and the labeling $d : X \to k$ is defined by $d(p) = c(p_{n-1})$. Relate to any explicit morphism $\varphi = (\varphi_0, \ldots, \varphi_{n-1}) : P \to Q$ of $\widetilde{\mathcal{P}}_k(n)$ the morphism $\varphi^\circ : P^\circ \to Q^\circ$ by $\varphi^\circ(p_0.p_1, \ldots) = (\varphi_0(p_0), \varphi_{1,p_0}(p_1), \ldots)$.

Conversely, relate to any object X of $\mathcal{S}_n(k)$ as above the object $X^+ = (P; \leq, c)$ of $\widetilde{\mathcal{P}}_k(n)$ where $(P; \leq)$ is the quotient-poset of $(X; \leq_0)$ and $c([x]_0) = ([x]_0; \leq_1, \ldots, \leq_{n-1}, d|_{[x_0]})^+$; we can suppose by induction that $c([x]_0)$ is an object of $\widetilde{\mathcal{P}}_k(n-1)$ if $n > 1$. Relate to any morphism $\psi : X \to Y$ of $\mathcal{S}_n(k)$ the explicit morphism $\psi^+ = (\varphi_0, \ldots, \varphi_{n-1}) : X^+ \to Y^+$ where $\varphi_0([x]_0) = [\psi(x)]_0$, $\varphi_{1,[x]_0}([z]_1) = [\psi(z)]_1$ for each $z \in [x]_0$ and so on. □

Note that the notion of an explicit morphism does not in fact depend on the number n because the explicit morphism $\varphi : P \to Q$ of $P, Q \in \widetilde{\mathcal{P}}_k(n)$ is uniquely extended to an explicit morphism of P, Q considered as objects of $\widetilde{\mathcal{P}}_k(n+1)$. Thus, we can consider the category $\widetilde{\mathcal{P}}_k(\omega)$ with the explicit morphism. Similarly, we can consider the category $\mathcal{S}_\omega(k) = \bigcup_n \mathcal{S}_n(k)$ of finite because $\mathcal{S}_n(k)$ may be considered as a subcategory of $\mathcal{S}_{n+1}(k)$ (just add the equality relation as $\leq_n$). In this way, we obtain the equivalence of categories $\widetilde{\mathcal{P}}_k(\omega)$ and $\mathcal{S}_\omega(k)$.

The full subcategories $\widetilde{\mathcal{F}}_k(\omega)$ and $\widetilde{\mathcal{T}}_k(\omega)$ of $\widetilde{\mathcal{P}}_k(\omega)$ are then equivalent to suitable full subcategories $\mathcal{U}_\omega(k)$ and $\mathcal{V}_\omega(k)$ of $\mathcal{S}_\omega(k)$ (e.g., the objects of $\mathcal{U}_\omega(k)$ are $(X; \leq_0 . \leq_1 \ldots, d)$ where $(X; \leq_0)$ is a forest, $([x]_0; \leq_1)$ is a forest for each $x \in X$, and so on).

We conclude this subsection with extending Proposition 4.7 to the structure $\widetilde{\mathcal{T}}_k^{\sqcup}(\omega)$. For this we need the ordinal $\xi = sup\{\omega_1, \omega_1^{\omega_1}, \omega_1^{(\omega_1^{\omega_1})}, \ldots\}$. According to the Cantor normal form, any non-zero ordinal $\alpha < \xi$ is uniquely representable in the form $\alpha = \omega_1^{\gamma_0} \cdot \alpha_0 + \cdots + \omega_1^{\gamma_l} \cdot \alpha_l$ for some $l < \omega$, $\alpha > \gamma_0 > \cdots > \gamma_l$ and non-zero ordinals $\alpha_0, \ldots, \alpha_l < \omega$). For $F \in \widetilde{\mathcal{F}}_2(\omega)$, let $\bar{F} \in \widetilde{\mathcal{F}}_2(\omega)$ be obtained from F by interchanging $\{0, 1\}$ in all the labels.

Definition 5.4 We define the sequence $\{T_\alpha\}_{\alpha < \xi}$ of trees in $\widetilde{\mathcal{T}}_2(\omega)$ by induction on α as follows:

(1) For $\alpha < \omega_1$, use the definition from the end of Subsection 4.2.
(2) For any non-zero ordinal $\gamma < \xi$, $T_{\omega_1^\gamma} = s(T_\gamma)$.
(3) For any limit uncountable ordinal $\lambda < \xi$ of countable cofinality, $T_\lambda = \bigsqcup_n T_{\alpha_n}$ where $\alpha_0 < \alpha_1 < \cdots$ and $\lambda = sup\{\alpha_0, \alpha_1, \ldots\}$.
(4) For any ordinal $\beta \geq \omega_1$, $T_{\beta+1} = 0 * (T_\beta \cup \bar{T}_\beta$.
(5) For all δ, γ such that $1 \leq \delta < \omega_1$ and $1 \leq \gamma < \xi$, $T_{\omega_1^\gamma(\delta+1)} = T_\gamma * \bar{T}_{\omega_1^\gamma \delta}$.
(6) For all β, γ such that $1 \leq \gamma < \xi$ and $\beta = \omega_1^{\gamma_1} \cdot \beta_1$ for some $\beta_1 > 0$ and $\gamma_1 > \gamma$, $T_{\beta+\omega_1^\gamma} = T_\gamma * (T_\beta \sqcup \bar{T}_\beta)$.
(7) For all δ, β, γ such that $1 \leq \delta < \omega_1$, $1 \leq \gamma < \xi$ and $\beta = \omega^{\gamma_1} \cdot \beta_1$ for some $\beta_1 > 0$ and $\gamma_1 > \gamma$, $T_{\beta+\omega_1^\gamma(\delta+1)} = T_\gamma * \bar{T}_{\beta+\omega_1^\gamma \delta}$.

The next assertion is the extension of Proposition 4.7. We omit the proof which is very similar to that of Proposition 8.29 in [Se12] which is a finitary version of this assertion.

Proposition 5.5 (1) *For all $\alpha < \beta < \xi$, T_α is correctly defined up to $\equiv_h$ (in particular, it does not depend on the choice of the ordinals α_n in Definition 5.4(3)), T_α and $\bar{T}_\alpha$ are $\leq_h$-incomparable elements of $\widetilde{\mathcal{T}}_2(\omega)$ that satisfy $T_\alpha \sqcup \bar{T}_\alpha <_h T_\beta$.*

(2) *Any $T \in \widetilde{\mathcal{T}}_2(\omega)$ is h-equivalent to precisely one of $T_\alpha, \bar{T}_\alpha$ for some $\alpha < \xi$.*

(3) *Any $T \in \widetilde{\mathcal{T}}_2^{\sqcup}(\omega)$ is h-equivalent to precisely one of $T_\alpha, \overline{T}_\alpha, T_\alpha \sqcup \overline{T}_\alpha, \bigsqcup_{\alpha<\lambda} T_\alpha$ where λ is a limit ordinal of countable cofinality.*

(4) *$T_\alpha \sqcup \overline{T}_\alpha \equiv_h T_{\alpha+1} \sqcap \overline{T}_{\alpha+1}$ and $\bigsqcup_{\alpha<\lambda} T_\alpha \equiv_h T_\lambda \sqcap \overline{T}_\lambda$ for any limit ordinal λ of countable cofinality, where $\sqcap$ is a binary operation of k-forests inducing the infimum operation in $\widetilde{\mathbb{F}}'_k(\omega)$.*

(5) *The ranks of $\widetilde{\mathbb{T}}_2(\omega)$ and of $\widetilde{\mathbb{T}}_2^{\sqcup}(\omega)$ coincide with ξ.*

5.2 Fine hierarchies of k-partitions

Here we define the TH of k-partitions and formulate its basic properties. The proofs are almost the same as those for the finitary case in [Se12].

Let X be a space. By an *ω-base* (cf. [Se08a, Se12]) in X we mean a sequence $\mathcal{L} = \{\mathcal{L}_n\}_{n<\omega}$ of bases such that $\mathcal{L}_n \cup \check{\mathcal{L}}_n \subseteq \mathcal{L}_{n+1}$ for each $n < \omega$. The main example of an ω-base is of course $\{\mathbf{\Sigma}^0_{n+1}\}_{n<\omega}$ but also other examples are interesting, in particular the finite shifts $\{\mathbf{\Sigma}^0_{m+n}\}_{n<\omega}$ for any fixed $1 \leq m < \omega$.

Let $P \in \widetilde{\mathcal{F}}_k(n)$ for some positive $n < \omega$. By a *P-family over $\mathcal{L}$* we mean a family $\{B_{p_0}, B_{p_0,p_1}, \dots, B_{p_0,\dots,p_{n-1}}\}$ where $p = (p_0, \dots, p_{n-1}) \in P^\circ$ (see the end of Subsection 5.1), $B_{p_0} \in \mathcal{L}_0, B_{p_0,p_1} \in \mathcal{L}_1, \dots, B_{p_0\dots,p_{n-1}} \in \mathcal{L}_{n-1}$, and the sets

$$\tilde{B}_{p_0} = B_{p_0} \setminus \bigcup\{B_r \mid r \leq_0 p\},\ \tilde{B}_{p_0,p_1} = B_{p_0,p_1} \setminus \bigcup\{B_r \mid r \leq_1 p\}, \dots$$

satisfy

$$\tilde{B}_{p_0} = \bigcup\{B_{p_0,p_1} \mid p_1 \in c(p_0)\},\ \tilde{B}_{p_0,p_1} = \bigcup\{B_{p_0,p_1,p_2} \mid p_1 \in c(p_0), p_2 \in c(p_1)\}, \dots.$$

To simplify notation, we often denote families just by $\{B_p\}$. Note that $d(p) = c(p_{n-1})$ is always in $\widetilde{\mathcal{F}}_k(0)$={0,. . . ,k-1},

$$\tilde{B}_{p_0} = \bigcup\{\tilde{B}_{p_0,p_1} \mid p_1 \in c(p_0)\},\ \tilde{B}_{p_0,p_1} = \bigcup\{\tilde{B}_{p_0,p_1,p_2} \mid p_1 \in c(p_0), p_2 \in c(p_1)\}, \dots$$

and that for $n = 1$ the P-families over $\mathcal{L}$ essentially coincide with the P-families of $\mathcal{L}_0$-sets in Subsection 4.3. Obviously, $\bigcup_{p_0} B_{p_0} = \bigcup_{p\in P^\circ} \tilde{B}_p$. We call a P-family $\{B_p\}$ over $\mathcal{L}$ *consistent* if $d(p) = d(q)$ whenever the components $\tilde{B}_p$ and $\tilde{B}_q$ have a nonempty intersection. Any such consistent P-family determines the k-partition $A : \bigcup_{p_0} B_{p_0} \to k$ where $A(x) = d(p)$ for some (equivalently, for any) $p \in P^\circ$ with $x \in \tilde{B}_p$; we also say in this case that A is defined by $\{B_p\}$. Note that this k-partition is determined by the defining P-family and it does not depend on the number n with $P \in \widetilde{\mathcal{F}}_k(n)$.

Let $\mathcal{L}^Y(P)$ be the set of k-partitions $A : Y \to k$ defined by some P-family over $\mathcal{L}$. In case $Y = X$ we omit the superscript X and call (temporarily) the family $\{\mathcal{L}(P)\}_{P\in\widetilde{\mathcal{F}}_k(\omega)}$ the FH of k-partitions over $\mathcal{L}$.

For $F \in \widetilde{\mathcal{F}}_k(\omega)$, by a *reduced F-family over $\mathcal{L}$* we mean a monotone F-family $\{B_p\}$ over $\mathcal{L}$ such that $B_{p_0} \cap B_{q_0} = \emptyset$ for all incomparable $p_0, q_0 \in F$, $B_{p_0,p_1} \cap B_{p_0,q_1} = \emptyset$ for all incomparable $p_1, q_1 \in c(p_0)$ and so on. Let $\mathcal{L}^Y_r(F)$ be the set of partial k-partitions defined by the reduced F-families $\{B_p\}$ over $\mathcal{L}$ such that $\bigcup_{p_0} B_{p_0} = Y$.

The next assertion is a a straightforward extension of Proposition 4.8 proved similarly to its finitary version in [Se12].

Proposition 5.6 (1) *If $A \in \mathcal{L}^Y(P)$ then $A|_Z \in \mathcal{L}^Z(P)$ for each $Z \subseteq Y$, $Z \in \mathcal{L}_0$.*

(2) *Any $A \in \mathcal{L}^Y(P)$ is defined by a monotone P-family $\{C_p\}$ (monotonicity means that $C_{q_0} \subseteq C_{p_0}$ for $q_0 \leq p_0$, $C_{p_0,q_1} \subseteq C_{p_0,p_1}$ for $q_1 \leq p_1$ and so on).*

(3) *Let $f : X_1 \to X$ be a morphism of ω-spaces and $A \in \mathcal{L}^Y(P)$ in X. Then $f^{-1}(A) \in \mathcal{L}^{f^{-1}(Y)}(P)$.*

(4) *If* $P \leq_h Q$ *then* $\mathcal{L}^Y(P) \subseteq \mathcal{L}^Y(Q)$.
(5) *The collection* $\{\mathcal{L}(P) \mid P \in \widetilde{\mathcal{F}}_k(\omega)\}$ *is well partially ordered by inclusion.*
(6) *For all* $F, G \in \widetilde{\mathcal{F}}'_k(\omega)$, $\mathcal{L}(F) \cap \mathcal{L}(G) = \mathcal{L}(F \sqcap G)$.
(7) *Any level* $\mathcal{L}(F)$, $F \in \widetilde{\mathcal{F}}_k(\omega)$, *of the FH is the intersection of finitely many "non-self-dual" levels* $\mathcal{L}(T)$, $T \in \widetilde{\mathcal{T}}_k(\omega)$.
(8) *Let* $\mathcal{L}_n$ *have the* ω*-reduction property for each* $n < \omega$, $Y \in \mathcal{L}_0$ *and* $F \in \widetilde{\mathcal{F}}_k(\omega)$. *Then* $\mathcal{L}^Y(F) = \mathcal{L}^Y_r(F)$, *in particular,* $\mathcal{L}(F) = \mathcal{L}_r(F)$.

The next result is an infinitary version of Proposition 8.19 in [Se12]. It extends the Hausdorff-Kuratowski theorem to all levels of the FH. We call the ω-base $\mathcal{L}$ is *interpolable* if, for each $n \geq 1$, $\mathcal{L}_n$ has the ω-reduction property and the Hausdorff-Kuratowski theorem holds for any non-zero level of $\mathcal{L}$. In particular, the base $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}(X)\}_{n<\omega}$ is interpolable in any quasi-Polish space.

Theorem 5.7 *If the* ω*-base* $\mathcal{L}$ *is interpolable then any non-empty constituent of the FH* $\{\mathcal{L}(x)\}_{x \in \widetilde{\mathbb{F}}_k(\omega)}$ *is a constituent of the FH* $\{\mathcal{L}(x)\}_{x \in \widetilde{\mathbb{T}}^{\sqcup}_k(\omega)}$.

Once we have the (hopefully) right definition of the FH of k-partitions, some properties of the DH are extended in a straightforward way. In particular, this applies to Propositions 4.11, 4.12 and Theorem 4.13. We give the corresponding formulations.

Proposition 5.8 *Let* $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}\}_{n<\omega}$ *and* $F \in \widetilde{\mathcal{F}}_k$. *Then* $\{\mathcal{L}(X, F)\}_X$ *is a family of* k*-partition classes.*

Proposition 5.9 *Let* X *be a* cb_0*-space,* $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+2}(X)\}_{n<\omega}$, *and* $T \in \widetilde{\mathcal{T}}_k$. *Then* $\mathcal{L}(X, T)$ *has a principal* $\mathcal{L}(T)$*-TR. If* X *is in addition zero-dimensional and* $\mathcal{M} = \{\mathbf{\Sigma}^0_{n+1}(X)\}_{n<\omega}$ *then* $\mathcal{M}(X, T)$ *has a principal* $\mathcal{M}(T)$*-TR.*

The next result extends (with essentially the same proof) Theorem 4.13 to the FH of k-partitions.

Theorem 5.10 *Let* X *be a* cb_0*-space,* $\delta : D \to X$ *an admissible representation of* X *(*$D \subseteq \mathcal{N}$*),* $A : X \to k$ *a* k*-partition of* X, $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}\}_{n<\omega}$, *and* $F \in \widetilde{\mathcal{F}}_k$. *Then* $A \in \mathcal{L}(X, T)$ *iff* $A \circ \delta \in \mathcal{L}(D, T)$.

Corollary 5.11 *Let* X *be a quasi-Polish space,* $\delta : \mathcal{N} \to X$ *an admissible TR of* X, $A : X \to k$ *a* k*-partition of* X, $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}\}_{n<\omega}$, *and* $F \in \widetilde{\mathcal{F}}_k$. *Then* $A \in \mathcal{L}(X, T)$ *iff* $A \circ \delta \in \mathcal{L}(\mathcal{N}, T)$.

For the DH of k-partitions, it was easy to demonstrate that it really extends the DH of sets (which coincides with the DH of 2-patitions by Proposition 4.9). For the FH of 2-partitions the same task is more complicated. The reason is that this hierarchy should generalize the Wadge hierarchy which is defined and relatively well understood only for the Baire space (and some other closely related spaces). Moreover, the most popular definition of this hierarchy is in terms of the Wadge reducibility rather than in set-theoretic terms (in fact, there is also set-theoretic definitions [Wad84, Lo83] but they are very indirect and hard to deal with). Nevertheless, we claim that the FH of 2-partitions in the Baire space coincides with the Wadge hierarchy. We discuss this (rather informally) in the next subsection.

5.3 FH of k-partitions and Wadge-like reducibilities

Let us briefly recall the definition of Wadge reducibility in the Baire space. In [Wad72, Wad84] W. Wadge (with a heavy use of the Martin determinacy theorem) proved that the structure $(\mathbf{\Delta}^1_1(\mathcal{N}); \leq_W)$ is semi-well-ordered (i.e., it is well-founded and for all $A, B \in \mathbf{B}$ we have $A \leq_W B$ or $\overline{B} \leq_W A$. He has also computed the rank ν of this structure which is a rather large ordinal.
In [VW76, St80] the following deep relation of the Wadge reducibility to the separation property was established: For any Borel set A which is non-self-dual (i.e., $A \not\leq_W \overline{A}$) exactly one of the principal ideals $\{X \mid X \leq_W A\}$, $\{X \mid X \leq_W \overline{A}\}$ has the separation property.
The mentioned results give rise to the *Wadge hierarchy* which is, by definition, the sequence $\{\mathbf{\Sigma}_\alpha\}_{\alpha<\nu}$ of all non-self-dual principal ideals of $(\mathbf{\Delta}^1_1(\mathcal{N}); \leq_W)$ that do not have the separation property and satisfy for all $\alpha < \beta < \nu$ the strict inclusion $\mathbf{\Sigma}_\alpha \subset \mathbf{\Delta}_\beta$. As usual, we set $\mathbf{\Pi}_\alpha = \{\overline{A} \mid A \in \mathbf{\Sigma}_\alpha\}$ and $\mathbf{\Delta}_\alpha = \mathbf{\Sigma}_\alpha \cap \mathbf{\Pi}_\alpha$.

Note that the constituents of the Wadge hierarchy are precisely the equivalence classes induced by $\leq_W$ on Borel subsets of the Baire space (i.e., the Wadge degrees).
As shown in [Wad84], $\mathbf{\Sigma}_\alpha = \mathbf{\Sigma}^{-1}_\alpha(\mathcal{N})$ for each $\alpha < \omega$, i.e. the DH over open sets is an initial segment of the Wadge hierarchy. In order to see how much finer is the Wadge hierarchy compared with the Borel hierarchy, we mention the equalities from [Wad84] relating both hierarchies: $\mathbf{\Sigma}_1 = \mathbf{\Sigma}^0_1(\mathcal{N})$, $\mathbf{\Sigma}_{\omega_1} = \mathbf{\Sigma}^0_2(\mathcal{N})$, $\mathbf{\Sigma}_{\omega_1^{\omega_1}} = \mathbf{\Sigma}^0_3$ and so on. Thus, the sets of finite Borel rank coincide with the sets of Wadge rank less than $\xi = sup\{\omega_1, \omega_1^{\omega_1}, \omega_1^{(\omega_1^{\omega_1})}, \ldots\}$. Note that ξ is the smallest solution of the ordinal equation $\omega_1^{\varkappa} = \varkappa$. Hence, we warn the reader not to mistake $\mathbf{\Sigma}_\alpha$ with $\mathbf{\Sigma}^0_\alpha$. To give the reader a first impression about the Wadge ordinal we note that the rank of the preorder $(\mathbf{\Delta}^0_\omega; \leq_W)$ is the ω_1-st solution of the ordinal equation $\omega_1^{\varkappa} = \varkappa$ [Wad84].
The structure of Wadge degrees is known to have the following properties: at the bottom (i.e., zero) level and at the limit levels of uncountable cofinality we have non-self-dual pairs of degrees; at the limit levels of countable cofinality we have self-dual degrees; at successor levels the self-dual degrees and non-sel-fdual pairs alternate. Remembering Proposition 5.5 we immediately see that the structure of Wadge degrees of finite Borel rank is isomorphic to the structure $\widetilde{\mathbb{T}}^{\sqcup}_2(\omega)$! This observation makes more plausible our claim that the FH of k-partitions extends the Wadge hierarchy.
To explain this more precisely, we note that it is possible to relate to any $F \in \widetilde{\mathcal{T}}^{\sqcup}_2(\omega)$ a k-partition A_F of the Baire space in such a way that A_F is Wadge complete in $\mathcal{L}(F)$ where $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}(\mathcal{N})\}_{n<\omega}$. Since the proof of this result is too technical for this paper, we postpone it to a subsequent publication and only note that very relevant particular cases are considered in [Se07, Se07a, Se11] (in fact, from these papers only the proof for the initial segment $\widetilde{\mathcal{T}}^{\sqcup}_k(2)$ is easily extracted, while for the general result one has to employ additional jump operators from [And06, MR09] in order to relate A_F to arbitrary $F \in \widetilde{\mathcal{T}}^{\sqcup}_k(\omega)$. In this way one obtains the following result showing, in particular, that the FH of sets really extends the Wadge hierarchy of sets of finite Borel rank:

Proposition 5.12 *For any $F \in \widetilde{\mathcal{T}}^{\sqcup}_k(\omega)$, A_F is Wadge complete in $\mathcal{L}(\mathcal{N}, F)$ where $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}(\mathcal{N})\}_{n<\omega}$ and, moreover, $F \leq_h G$ iff $\mathcal{L}(\mathcal{N}, F) \subseteq \mathcal{L}(\mathcal{N}, G)$.*
In particular, for any $\alpha < \xi$ we have $\mathcal{L}(\mathcal{N}, T_\alpha) = \mathbf{\Sigma}_\alpha$ where $T_\alpha \in \widetilde{\mathcal{T}}_2(\omega)$ is the tree from Proposition 5.5.

Our definition of the FH of k-partitions is thus a natural extension of the Wadge hierarchy to arbitrary spaces. Interestingly, for the important particular case of quasi-Polish spaces this hierarchy is naturally induced from the Wadge hierarchy of k-partitions of the Baire space via admissible representations. This follows immediately from Corollary 5.11.

Theorem 5.13 *Let X be a quasi-Polish space, $\mathcal{L} = \{\mathbf{\Sigma}^0_{n+1}(\mathcal{N})\}_{n<\omega}$, $\mathcal{M} = \{\mathbf{\Sigma}^0_{n+1}(X)\}_{n<\omega}$, and $F \in \widetilde{\mathcal{T}}^{\sqcup}_k(\omega)$. Then $A \in \mathcal{M}(X, F)$ iff $A \circ \delta \in \mathcal{L}(\mathcal{N}, F)$ where δ is some (equivalently, any) admissible TR of X.*

As is well known, the structure of Wadge degrees in many natural non-zero-dimensional cb_0-spaces is very complicated (see e.g. [MSS12] and references therein) so it seems hopeless to understand these structures well enough. Nevertheless, if we slightly weaken the notion of Wadge reducibility by extending the class of reducing functions (this is similar to the relativization process in Computability Theory) we obtain natural versions of Wadge reducibility which behave similar to the classical one in many natural spaces. This also applies to reducibilities of k-partition. We illustrate this with the following assertion:

Proposition 5.14 *Let X be a quasi-Polish space such that $dim(X) \neq \infty$. Then the structure of $\mathbf{\Sigma}^0_3$-degrees of k-partitions of X of finite Borel rank is isomorphic to $\widetilde{\mathcal{T}}^{\sqcup}_k(\omega)$.*

Proof Hint. By Proposition 3.1, there is a $\mathbf{\Sigma}^0_3$-isomorphism f between X and $\mathcal{N}$. Clearly, f induces an isomorphism of the quotient-structures of $(k^X; \leq_{\mathbf{\Sigma}^0_3})$ and $(k^{\mathcal{N}}; \leq_{\mathbf{\Sigma}^0_3})$ which preserves the initial segments of k-partitions of finite Borel rank. Therefore, it suffices to prove the assertion for $X = \mathcal{N}$. But this is just the $\mathbf{\Sigma}^0_3$-relativization of Proposition 5.12. Note that similar relativizations are employed in [MR09]. □

6 Conclusion

The results of this paper suggest that DST in cb_0-spaces (or at least in some rich classes of cb_0-spaces) resembles in many respects the classical DST in Polish spaces. Also, the methods of classical DST seem also to work well in this context, although a more systematic treatment of DST in cb_0-spaces is desirable. In particular, the classical theory of equivalence relations and descriptive theory of functions on cb_0-spaces (more complicated than the k-partitions considered here) seem to be interesting.
Of course, many details on the FH of k-partitions in cb_0-spaces should be elaborated much more carefully than in this paper. In fact, even the structure of Wadge degrees of Borel k-partitions of the Baire space should be described much more carefully; we plan to do this in subsequent publications (of course one cannot expect to fulfill this task in a short single paper because even much easier case of sets is technically very involved [Wad84, VW76], and the game-theoretic technique does not work in non-zero-dimensional spaces). The results of this paper suggest that also the FH of k-partitions in arbitrary quasi-Polish spaces, and even in more general cb_0-spaces, is tractable.
A special challenge is the systematic development of DST in non-countably based spaces, in particular, in reasonable rich enough classes of qcb_0-spaces. This task could require also new methods compared with the case of cb_0-spaces.
Another interesting direction is the development of effective DST in effective (in some reasonable sense) spaces. As is well known from Computability Theory, this task is highly non-trivial even for "simple" spaces like the Baire space. For topologically more complicated spaces this direction is still widely open, although it seems of principal importance, in particular for Computable Analysis.

References

[And06] A. Andretta, More on Wadge determinacy, Ann. Pure Appl. Logic 144, 2006, no. 1-3, 2–32.

[AM03] A. Andretta and D. A. Martin, Borel-Wadge degrees, Fund. Math. 177, 2003, no. 2, 175–192.

[Br13] M. de Brecht. Quasi-Polish spaces. *Annals of pure and applied logic*, 164 (2013), 356–381.

[BY09] M. de Brecht and A. Yamamoto. $\mathbf{\Sigma}^0_\alpha$-admissible representations (Extended Abstract). Proc. of 6-th Int. Conf. on Computability and Complexity in Analysis, 2009.

[Du01] Duparc J. Wadge hierarchy and Veblen hierarchy. Part I: Borel Sets of Finite Rank. Journal of Symbolic Logic, 66(1), 2001, 56-86.

[En89] R. Engelking. *General Topology.* Heldermann, Berlin, 1989.

[FHK11] S.D. Friedman, T. Hyttinen, V. Kulikov. Generalized descriptive set theory and classification theory. CRM preprint no 999, 2011.

[GH80] G. Giertz, K.H. Hoffmann, K. Keimel, J.D. Lawson, M.W. Mislove, D.S. Scott. *A Compendium of Continuous Lattices.* Berlin, Springer, 1980.

[Her93] P. Hertling. Topologische Komplexitätsgrade von Funktionen mit endlichem Bild. Informatik-Berichte 152, 34 pages, Fernuniversität Hagen, December 1993.

[Her96] P. Hertling. *Unstetigkeitsgrade von Funktionen in der effektiven Analysis.* PhD thesis, Fachbereich Informatik, FernUniversität Hagen, 1996.

[JR82] J.E. Jayne and C.A. Rogers, First level Borel functions and isomorphisms, J. Math. Pures Appl. (9) **61** (1982), no. 2, 177–205.

[Ke83] A.S. Kechris. Suslin cardinals, k-Suslin sets and the scale property in the hyperprojective hierarchy. The Cabal Seminar, v. 1: Games, Scales and Suslin Cardinals, Eds. A.S. Kechris, B. Löwe, J.R. Steel, Lecture Notes in Logic, 31, 2008, p. 314–332. (Reprinted from Lecture Notes in mathematica, No 1019, Berlin, Springer, 1983).

[Ke95] A.S. Kechris. *Classical Descriptive Set Theory.* Springer, New York, 1995.

[Kru72] J.B. Kruskal. The theory of well-quasi-ordering: a frequently discovered concept. *J. Combinatorics Th.(A)*, 13 (1972), 297–305.

[KW85] C. Kreitz and K. Weihrauch (1985). Theory of representations. *Theoretical Computer Science*, 38:35–53.

[Lo83] A. Louveau. Some results in the Wadge hierarchy of Borel sets. *Lec. Notes in Math.*, No 1019 (1983), p.28–55.

[MR09] L. Motto Ros, Borel-amenable reducibilities for sets of reals, J. Symbolic Logic **74** (2009), no. 1, 27–49.

[MS10] L. Motto Ros and B. Semmes, A new proof of a theorem of Jayne and Rogers, Real Anal. Exchange **35** (2010), no. 1, 195–203.

[MSS12] L. Motto Ros, P. Schlicht and V. Selivanov. Wadge-like reducibilities on arbitrary quasi-Polish spaces. To appear in Mathematical Structures in Computer Science, arXiv:1204.5338 v1 [mathLO] 24 Apr 2012.

[P12] A. Pauly. A new introduction to the theory of represented spaces, arXiv:1204.3763, 2012.

[PdB13] A. Pauly and M. de Brecht. Towards synthetic descriptives set theory: An instantiation with represented spaces, arXiv:1307.1850, 2013.

[Sch02] M. Schröder. *Extended admissibility.* Theoretical Computer Science 284 (2002), 519–538.

[Sch03] M. Schröder. *Admissible representations for continuous computations.* PhD thesis, Fachbereich Informatik, FernUniversität Hagen, 2003.

[ScSe13] M. Schröder, V. Selivanov. Some hierarchies of qcb_0-spaces. *Logical Methods in Computer Science*, to appear in *Mathematical Structures in Coputer Science*. arXiv:1304.1647

[ScSe14] M. Schröder, V. Selivanov. Hyperprojective hierarchy of qcb_0-spaces. To appear in LNCS-volume of CiE-2014.

[Se04] V.L. Selivanov. Difference hierarchy in φ-spaces. *Algebra and Logic*, 43, N 4 (2004), 238–248.

[Se05] V.L. Selivanov. Variations on the Wadge reducibility. *Siberian Advances in Mathematics*, 15, N 3 (2005), 44–80.

[Se05a] V.L. Selivanov. Hierarchies in φ-spaces and applications. *Math. Logic Quarterly*, 51, N 1 (2005), 45–61.

[Se06] V.L. Selivanov. Towards a descriptive set theory for domain-like structures. *Theoretical Computer Science*, 365 (2006), 258–282.

[Se07] V.L. Selivanov. The quotient algebra of labeled forests modulo h-equivalence. *Algebra and Logic*, 46, N 2 (2007), 120–133.

[Se07a] V.L. Selivanov. Hierarchies of $\mathbf{\Delta}^0_2$-measurable k-partitions. *Math. Logic Quarterly*, 53 (2007), 446–461.

[Se08] V.L. Selivanov. On the difference hierarchy in countably based T_0-spaces. *Electronic Notes in Theoretical Computer Science*, V. 221 (2008), 257-269.

[Se08a] V.L. Selivanov. Fine hierarchies and m-reducibilities in theoretical computer science. *Theoretical Computer Science*, 405 (2008), 116–163.

[Se10] V.L. Selivanov. On the Wadge reducibility of k-partitions. Journal of Logic and Algebraic Programming, 79, No 1, 2010, 92–102.

[Se11] V.L. Selivanov. A fine nierarchy of ω-regular k-partitions. B. Löwe et.al. (Eds.): CiE 2011, LNCS 6735, pp. 260–269. Springer, Heidelberg (2011).

[Se12] V.L. Selivanov. Fine hierarchies via Piestley duality. Annals of Pure and Applied Logic, 163 (2012) 1075-1107

[Se13] V.L. Selivanov. Total representations. *Logical Methods in Computer Science*, 9(2) (2013), p. 1–30.

[SR07] J. Saint Raymond. Preservation of the Borel classes under countable-complete-covering mappings. Topology and its Applications, 154 (2007), 1711–1725.

[St80] J. Steel. Determinateness and the separation property. *J. Symbol. Logic,* 45 (1980), p.143–146.

[VW76] R. Van Wesep. Wadge degrees and descriptive set theory. *Lec. Notes in Math.,* No 689 (1976), p. 151–170.

[Wad72] W. Wadge. Degrees of complexity of subsets of the Baire space. *Notices AMS*, 1972, A-714.

[Wad84] W. Wadge. *Reducibility and determinateness in the Baire space.* PhD thesis, University of California, Berkely, 1984.

[We00] K. Weihrauch. *Computable Analysis.* Berlin, Springer, 2000.